


Norges teknisk-naturvitenskapelige universitet - NTNU
Fakultet for naturvitenskap og teknologi
Institutt for fysikk



HOVEDOPPGAVE FOR

STUD. TECHN. GEIR OVE MYHR

Oppgaven gitt: 01.02.2004
Besvarelsen levert: 27.06.2004

FAGOMRÅDE: **TEORETISK FYSIKK**

Norsk tittel: *“Mål på sammenfiltring i kvantemekanikk”*

Engelsk tittel: *“Measures of entanglement in quantum mechanics”*

Hovedoppgaven er utført ved Institutt for fysikk, Fakultet for naturvitenskap og teknologi,
under veiledning av professor Jan Myrheim og stipendiat Stein Olav Skrøvseth.

Trondheim, 27. juni 2004,

Jan Myrheim
ansvarlig faglærer
professor ved Institutt for fysikk

Abstract

I give an overview of some of the most used measures of entanglement. To make the presentation self-contained, a number of concepts from quantum information theory are first explained. Then the structure of bipartite entanglement is studied qualitatively, before a number of bipartite entanglement measures are described, both for pure and mixed states. Results from the study of multipartite systems and continuous variable systems are briefly discussed.

Preface

“What’s the name again of this quantum effect when one particle on one side of the universe magically affects a particle on the other side of the universe?”, I asked while walking out the door to the corridor. I was on my way to a group of teenagers who were waiting to hear how it is to study physics at the university. In case someone asked why I found physics so exciting I could always mention this quantum, non-locality, spooky action-at-a-distance thing. If I only remembered the name of it... “That’d be *entanglement*”, a graduate student next-door suggested, using the English term. Yes, that was the word lost in the back of my mind. I had actually never heard a Norwegian name for it.

This happened one of the last days of October 2003. About a month later it was decided that my master’s thesis would be about this very action-at-a-distance thing, or more precisely, about how to quantify entanglement. From hardly remembering the name of the game to understanding how an appropriate measure needs to behave has been a long and interesting journey. In writing this text I have tried to keep in mind my own state of knowledge as of October 2003. With that I hope that any 5th year physics student will find it sufficiently understandable and self-contained as to be able to grasp the main points. Whether I have succeeded or not is for others to decide.

This thesis is the final work leading up to the degree of Master of Science in Applied Physics and Mathematics at the Norwegian University of Science and Technology (NTNU). It was written during the 2004 spring term.

I would like to thank Jan Myrheim and Stein Olav Skrøvseth for taking the extra effort to find a topic in quantum information theory for my thesis, and for supervising it. I would also like to thank Viktor H. Havik and Henrik Tollefsen for interesting discussions about entanglement.

Contents

Abstract	v
Preface	vii
Contents	ix
1 Introduction	1
2 Preliminaries	3
2.1 The qubit	3
2.2 Entangled states	5
2.3 Density operators and mixed states	7
2.4 The Schmidt decomposition	10
2.5 Generalised measurements	11
2.6 Transformation of states and LOCC	13
2.6.1 Local Operations and Classical Communication	13
2.6.2 Transformation of a Bell state	14
2.6.3 Concentrating partial pure state entanglement	14
2.6.4 Distilling entanglement from mixed states	16
3 Characterisation of bipartite entanglement	21
3.1 Pure states	21
3.2 Mixed states	22
3.2.1 Separable states	22
3.2.2 Free and bound entanglement	23
4 Measures of bipartite entanglement	25
4.1 Pure states	25
4.1.1 von Neumann entropy	26
4.1.2 Reduced von Neumann entropy – entropy of entanglement	26
4.1.3 Entanglement cost and distillable entanglement	27
4.1.4 The uniqueness theorem for measures of entanglement	28
4.2 Mixed states	29
4.2.1 Entanglement cost and distillable entanglement for mixed states	29
4.2.2 Axioms for abstract measures	30

4.2.3	E_D and E_C as extreme measures	32
4.2.4	Relative entropy of entanglement and other distance based measures .	33
4.2.5	Entanglement of formation	35
4.2.6	Negativity	37
4.2.7	Squashed entanglement	39
4.2.8	Other entanglement measures	40
5	Beyond bipartite entanglement and finite dimension	43
5.1	Multipartite systems	43
5.1.1	Separability	44
5.1.2	Measures of entanglement	44
5.2	Continuous variables	45
A	Expressions for the quantum relative entropy	47
	References	49
	Index	57

Chapter 1

Introduction

Quantum entanglement was first viewed as a curiosity when it was pointed out in the attempt by Einstein, Podolsky and Rosen [EPR35] to show that quantum mechanics could not be a complete theory. When John Bell many years later published his now well-known theorem [Bel64], it was hardly noticed in the scientific community. Bell had showed that for a theory based on local hidden variables, certain correlations were upper bounded, a result today known as the Bell inequalities. Quantum mechanics predicted that the inequalities would be violated, thus giving rise to a way of testing whether the predictions of quantum mechanics or the assumptions of Einstein *et al.* were correct. Experiments have later agreed with quantum mechanics [ADR82], although there are still critics arguing that there are loopholes in the experimental assumptions [TH02, San04].

Entanglement has since been regarded as a real – albeit strange – phenomenon of quantum mechanics. It was widely regarded as being the same as violation of some Bell inequality. However, Werner [Wer89] showed that there exist mixed quantum states that allow for a local hidden-variable theory, but nevertheless are entangled. This led to new criteria to decide if a state was separable [Per96, HHH96].

Parallel to this, the view of entanglement had gradually changed. From being regarded as a curiosity, it was now seen as an information processing and communication resource that could be used for performing tasks that would be impossible without it. Among the possibilities introduced were quantum cryptography [Eke91], dense coding [BW92], teleportation of a quantum state [BBC⁺93] and exponential speed-up of certain computational tasks [Sho94, Sho97, Gro96, Gro97]. Because entanglement was now viewed as a resource, it was natural to try to *quantify* the entanglement in quantum states.

After the first papers on the quantification of entanglement [SM95, BBP⁺96, BBPS96, BDSW96], the subject has grown into a whole field of research. The aim of this work is to describe the entanglement measures that have come out of this. The main focus is on entanglement between two parties with finite dimensional Hilbert spaces, as this is the situation where most of the theory is known. Extensions to entanglement between more than two parties and entanglement with continuous variables are also briefly discussed.

In the course of writing this, several reviews of entanglement and quantum information theory have been of great help. The textbook by Nielsen and Chuang [NC00] gives a nice introduction to the prerequisites to understand entanglement theory, but contains little on

entanglement itself. John Preskill's lecture notes [Pre98] explain much of the same, but has a whole chapter devoted to entanglement. The tutorial by Bruß [Bru02] gives a short introduction to most aspects of entanglement, including some entanglement measures. Both the lecture notes by Eisert [Eis03] and the review article by Keyl [Key02] include a chapter on entanglement measures, but the latter is rather technical in nature. The review article by M. Horodecki about entanglement measures [Hor01] summarises the most of the field in an elegant way, but it requires much prior knowledge from the reader, and treats almost exclusively two-party entanglement.

The rest of this work is organised as follows. Chapter 2 gives an introduction to the concepts needed in the following chapters. A reader with a basic knowledge in quantum information theory may skip large portions of this chapter. Chapter 3 describes two-party entanglement qualitatively, preparing the ground for chapter 4 which describes the ways to quantify two-party entanglement. Finally, in chapter 5 it is briefly indicated how the notions from the previous chapters may be extended to systems of continuous variables and to multi-party systems.

Unless stated otherwise all logarithms are taken base-2.

Chapter 2

Preliminaries

Before we can understand how to quantify entanglement, we need to have some concepts and procedures clear. In this chapter we will have a first look at qubits and entangled states, get used to density operators and see how quantum states can be transformed into others.

2.1 The qubit

The simplest nontrivial quantum mechanical system is a two-state system which can be described by a vector in two-dimensional complex Hilbert space. Such a system is called a *qubit*. The favourite qubit of many physicists is the spin of a spin-1/2 particle, for instance an electron. We will use spins of such particles to illustrate entanglement.

If we measure the z -component of the spin, the result is either up or down. This measurement corresponds to the observable operator which we call Z . After the measurement the state of the particle is an eigenstate of the observable. We denote these states by $|\uparrow_z\rangle$ and $|\downarrow_z\rangle$, for spin parallel to and antiparallel to the z -axis. They are mutually orthogonal, and we take them as the unit basis vectors of the spin Hilbert space. In addition to the eigenstates, the particle may be in any superposition of the eigenstates,

$$|\psi\rangle = \alpha |\uparrow_z\rangle + \beta |\downarrow_z\rangle \quad (2.1)$$

such that $|\alpha|^2 + |\beta|^2 = 1$. When the spin along the z -axis is measured, we obtain $+1$ with a probability of $|\alpha|^2$ and -1 with probability $|\beta|^2$ (where the spin is in units of $\hbar/2$). After the measurement the state collapses into the corresponding eigenstate.

As the overall phase of a quantum state does not have any physical significance, only the phase difference between α and β is important, and for convenience we often choose α to be real. In order to also include the property $|\alpha|^2 + |\beta|^2 = 1$, we can write the spin state (2.1) of the particle as

$$|\psi\rangle = \cos \frac{\theta}{2} |\uparrow_z\rangle + e^{i\phi} \sin \frac{\theta}{2} |\downarrow_z\rangle. \quad (2.2)$$

This gives a nice and intuitive picture of the state space of the spin-1/2 particle, and thereby any qubit. All possible states are covered when the parameters are restricted to $0 \leq \phi \leq 2\pi$ and $0 \leq \theta \leq \pi$. Thus, any state corresponds to a point on the unit sphere, with the

azimuthal angle given by ϕ and the polar angle given by θ . Such a sphere is called a *Bloch sphere*, and is a very useful tool for visualising states of a qubit. A Bloch sphere with some states indicated is shown in figure 2.1. Note that for $\theta = \pi$, the phase of (2.2) changes as ϕ changes, but the state is still the same. This is reflected on the Bloch sphere, as the azimuthal angle is irrelevant at the poles.




Figure 2.1: The Bloch sphere of the Hilbert space spanned by $|\uparrow_z\rangle$ and $|\downarrow_z\rangle$.

We may choose another set of orthonormal basis vectors. For instance

$$|\uparrow_x\rangle = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle + |\downarrow_z\rangle) \quad |\downarrow_x\rangle = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle - |\downarrow_z\rangle). \quad (2.3)$$

These are the eigenvectors of the X -operator, and they are the resulting states if we measure the spin along the x -axis. Likewise, we may measure the spin along any axis, and find a corresponding set of eigenvectors. If we denote the basis vectors

$$|\uparrow_z\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad |\downarrow_z\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (2.4)$$

then the operators for the spin along the x , y and z axes are nothing more than the Pauli matrices

$$X \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad Y \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad Z \equiv \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.5)$$

For the spin along any other unit vector, say $a\vec{x} + b\vec{y} + c\vec{z}$ where $a^2 + b^2 + c^2 = 1$, the spin operator is $aX + bY + cZ$. All operators of this form have eigenvalues ± 1 corresponding to spin parallel and antiparallel to the vector.

As already mentioned, other systems are equally good qubits as a spin-1/2 particle. Among the more popular ones are the polarisation of a photon and the energy levels of a two-level system. They can be represented by a Bloch sphere just as well as the spin of an electron can. The advantage of thinking about a spin-1/2 particle is that the vectors on the Bloch sphere corresponds to actual spin directions in real space. Thus, a qubit in a state close to the y -axis on the Bloch sphere, has, when the qubit is a spin-1/2 particle, a spin pointing in a direction close to the y -axis.

In general the particular implementation of the qubit is not important for our purpose, so we simply denote the basis vectors $|0\rangle$ and $|1\rangle$, which in the case of a spin-1/2 particle are typically $|\uparrow_z\rangle$ and $|\downarrow_z\rangle$, respectively. The superpositions of these states corresponding to the spin along the x -axis, are often denoted $|+\rangle$ and $|-\rangle$, according to the sign in the superposition.

$$|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \quad |-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \quad (2.6)$$

Three qubit operators are of special importance, namely the unitary Pauli operators of (2.5). In the context of general qubits, they can be written as

$$X \equiv |1\rangle\langle 0| + |0\rangle\langle 1|, \quad (2.7a)$$

$$Y \equiv i|1\rangle\langle 0| - i|0\rangle\langle 1|, \quad (2.7b)$$

$$Z \equiv |0\rangle\langle 0| - |1\rangle\langle 1|. \quad (2.7c)$$

X is also called the *bit flip operator* as it turns a $|0\rangle$ into a $|1\rangle$ and vice versa. Z is called the *phase flip operator* as it switches between the states $\alpha|0\rangle + \beta|1\rangle$ and $\alpha|0\rangle - \beta|1\rangle$. The Pauli operators are defined with respect to a particular basis. For instance, in the $\{|+\rangle, |-\rangle\}$ basis, Z takes the role of the bit flip operator.

The name *qubit* comes from *quantum bit* and it plays the same role in quantum information theory as the bit does in classical information theory. The bit is a system which can be in one of two states, usually denoted 0 and 1. Thus, each bit is capable of storing enough information to answer exactly one yes/no question. As we have just seen, the qubit is a much richer structure. In addition to being capable of being in the states $|0\rangle$ and $|1\rangle$, just as a bit, it can be in a continuum of superpositions between those states, given by two parameters. Still, only a single bit of information can be read from a single qubit. When the spin of a spin-1/2 particle is measured along an axis the only possible results are parallel and antiparallel to the chosen axis. After the measurement the state is changed, and from the result of the measurement we know what it is. No more information about the original state can be retrieved.

2.2 Entangled states

From the simplest quantum systems whatsoever, we now turn to look at the simplest *composite* quantum systems. A composite system is a system which consists of two or more parts, and the simplest one is a system consisting of two qubits. We call the two systems A and B . In communication and entanglement theory the convention is to think of the

subsystems as being in the possession of two well separated observers, called Alice and Bob. Any state of each of the two systems can be written as

$$|\psi\rangle_A = \alpha|0\rangle_A + \beta|1\rangle_A \quad |\phi\rangle_B = \gamma|0\rangle_B + \delta|1\rangle_B \quad (2.8)$$

with $|\alpha|^2 + |\beta|^2 = 1$ and $|\gamma|^2 + |\delta|^2 = 1$. The composite state of the two systems is then simply the tensor product (or direct product) of the two states.

$$|\Psi_{\text{prod}}\rangle = |\psi\rangle_A \otimes |\phi\rangle_B \quad (2.9)$$

Such a state is called a *product state*, but product states are not the only physically realisable states. If we let the two systems interact with each other, any superposition of product states is realisable. Hence, a general composite state can be written as

$$|\Psi\rangle = \sum_{ij} \alpha_{ij} |\psi_i\rangle_A \otimes |\phi_j\rangle_B \quad (2.10)$$

where $\sum |\alpha_{ij}|^2 = 1$ and the sets $\{|\psi_i\rangle\}$ and $\{|\phi_j\rangle\}$ are orthonormal bases for the two subsystems. Any composite state that is not a product state is called an *entangled state*.

A composite quantum state consisting of two parts only, is called a *bipartite state*, as opposed to *multipartite states* which consist of more than two parts. For bipartite qubit states, four entangled states play a major role, namely the singlet state

$$|\Psi^-\rangle \equiv \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle) \quad (2.11a)$$

and the three triplet states

$$|\Psi^+\rangle \equiv \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle) \quad (2.11b)$$

$$|\Phi^-\rangle \equiv \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle) \quad (2.11c)$$

$$|\Phi^+\rangle \equiv \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \quad (2.11d)$$

where we have used $|ij\rangle$ as a shorthand notation for $|i\rangle \otimes |j\rangle$. They are called *Bell states* or *EPR pairs*. Together they form an orthonormal basis for the state space of two qubits, called the *Bell basis*. The Bell states are maximally entangled and one can be converted into another by applying a unitary transform locally on any one of the subsystems.

Note that if we measure the state of one qubit in a Bell state (that is, measure the Z operator which has eigenvalues ± 1), we immediately know the state of the other particle. In the singlet Bell state, a measurement of qubit A will yield one of the eigenstates $|0\rangle$ and $|1\rangle$, each with a probability of $1/2$. These results leave qubit B in state $|1\rangle$ or $|0\rangle$, respectively.

For a single qubit we could always change to another basis where the outcome of a Z measurement would be given. For a spin-1/2 particle this means that the spin is always pointing in *some* direction, even though the state will show up as a superposition in a basis where the state is not one of the basis states. If the particle is entangled with another particle, though, the direction of the spin of that particle alone is not well defined. Actually, for particles in one of the Bell states, the probability for measuring the spin of the particle to “up” (while ignoring the other particle) is $1/2$ for *any* direction.

2.3 Density operators and mixed states

We want to be able to describe the state of a subsystem that is entangled with another subsystem to which we do not have access. State vectors cannot be used for this purpose, and we need another representation of quantum states, namely *density operators*. This is not only useful for describing “locally” a subsystem which is part of an entangled system for which we know the state. The formalism is also necessary for describing quantum mechanical experiments – where noise is inevitable – and doing quantum statistical mechanics. In the laboratory it is impossible to isolate the quantum systems under study completely. The systems become entangled with the environment through unwanted, but unavoidable, interactions. Nevertheless, it is necessary to describe the system without taking the environment into account.

In the density operator formalism, we describe quantum states by operators on the system’s Hilbert space instead of unit vectors on it. For any quantum state vector $|\psi\rangle$, the corresponding density operator is the projection operator $|\psi\rangle\langle\psi|$. As linear operators may be – and often is – represented by matrices, density operators are also called *density matrices*.

So far we have only represented the state in a new way, using density operators instead of vectors. But what if we do not know exactly in what state our system is? Imagine, for example, that we have a machine in our lab which outputs particles in state $|\psi\rangle$ (think of it as some component of a spin). But once in a while events beyond our control and knowledge (a spike in the electric grid, the turning on of an electromagnet in a neighbouring lab, etc.) make the machine malfunction slightly producing states which are close to, but not exactly $|\psi\rangle$. Let us for simplicity say that the state $|\psi'\rangle$ is always created when the machine malfunctions, and that the probability for that to happen to any given particle is p . Given this uncertainty, the state cannot be described by a state vector. Still, in terms of density matrices it is described as

$$\rho = (1 - p) |\psi\rangle\langle\psi| + p |\psi'\rangle\langle\psi'|.$$

A state that may be represented by a unit vector is called a *pure state*. For a pure state we have *maximal knowledge* of the state. Other states are *convex combinations*¹ of pure states, and are called *mixed states*. To check if a given density operator, ρ , represents a pure state, it is sufficient to check if $\rho = \rho^2$, which holds for all pure states and no mixed states.

The density operator does not tell us from what pure states the state was prepared. For example, we can have a state $|0\rangle$ or $|1\rangle$, each with probability $1/2$. The density operator is then $\frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|) = \frac{1}{2}\mathbb{I}$. The same density operator can be made by mixing the states $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ and $|-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$ in equal amounts, or the three states $|0\rangle$, $|+\rangle$ and $\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle$ in amounts $p_1 = \frac{1}{2} - \frac{\sqrt{3}}{6}$, $p_2 = \frac{\sqrt{3}}{3+\sqrt{3}}$ and $p_3 = 1 - \frac{\sqrt{3}}{3}$, respectively. We call these different collections of pure states with corresponding probabilities, $\{p_i, |\psi_i\rangle\}$, *ensembles*. Sometimes it is also called a *realisation* and we say that an ensemble *realises* a

¹A convex combination of elements $x_1, x_2, \dots, x_N$ is an element which can be written $\sum_i p_i x_i$ for $p_i \geq 0$ and $\sum_i p_i = 1$. A convex combination of two points is a point on the straight line connecting them.

mixed state. Hence, when we describe a state from an ensemble by its density operator, we discard the information about which ensemble the mixed state was made from. The density operator still describes the mixed state as well as can be done, as states from different ensembles having the same density operator are experimentally indistinguishable.

This indistinguishability makes the density operator a perhaps more intuitive representation of a state than the state vector. For example, a state vector has an arbitrary overall phase factor, so two vectors differing by a phase factor represent the same state. Still, when the vector is put together with its dual vector to form a pure state density operator, the phase factor vanishes because of the complex conjugation. Hence, all state vectors representing the same state are represented by the same density operator.

To complete our discussion about density operators, we should take note of some of their properties. For a more thorough discussion, see e.g. [NC00]. Density operators are (i) *positive*², meaning that for any vector $|v\rangle$, $\langle v|\rho|v\rangle \geq 0$, or equivalently that it is Hermitian with nonnegative eigenvalues. Also, (ii) the trace of a density operator is unity, $\text{tr}(\rho) = 1$. It turns out that any operator satisfying these two criteria can be realised by a pure state ensemble. Hence, we may take these criteria as the defining properties of a density operator.

We will now see how the density operators are used to describe individual parts of an entangled system. In fact, we will see that even when the composite system is in a pure state, if the subsystems are entangled, the individual subsystems are in mixed states.

Consider a bipartite composite system of two qubits, shared between Alice and Bob. The qubits are entangled, and the system is in the state

$$|\psi\rangle = \alpha|00\rangle + \beta|11\rangle. \quad (2.12)$$

Now, imagine that Alice loses contact with Bob, and wants to describe her system without any reference to the qubit in Bob's lab. Whatever Bob happens to be doing to his qubit should not have observable consequences in Alice's lab. If it had, it could have been used to perform superluminal communication in some reference frame which would wreak havoc in physics by allowing consequences to happen before the cause³. One of the things Bob might choose to do with his system, is to measure Z in the $\{|0\rangle, |1\rangle\}$ basis. If he does, he will get $|0\rangle$ or $|1\rangle$ with probability $|\alpha|^2$ or $|\beta|^2$, respectively. This leaves Alice's qubit in the same state as the one Bob measured, but as she wouldn't know the result, the state of her qubit would be a mixture of $|0\rangle$ and $|1\rangle$, namely

$$\rho_A = |\alpha|^2|0\rangle\langle 0| + |\beta|^2|1\rangle\langle 1| \quad (2.13)$$

This mixture correctly predicts the probabilities for all measurements that can be performed locally by Alice, regardless of what Bob really does to his system.

Formally, ignoring some degrees of freedom in a system (like the ones corresponding to a particle out of reach) is done by *tracing out* the relevant degrees of freedom from the density

²Positive operators here refer to *positive semidefinite* operators. *Positive definite* operators, on the other hand, satisfy $\langle v|\rho|v\rangle > 0$ for any $|v\rangle$ and have only positive eigenvalues.

³This is of course in itself no compelling reason. Even John Bell was afraid that entanglement might do away with relativity [Whi98]. The real reason is that quantum mechanics tells us that the probabilities for outcomes of local measurements on one system do not change when an arbitrary operation is performed on the other system.

operator. This is also called taking the *partial trace* with respect to the degrees of freedom that are to be traced out. Any bipartite state can be written as

$$\rho = \sum_{ijkl} c_{ijkl} |a_i\rangle \langle a_j| \otimes |b_k\rangle \langle b_l|$$

and the partial trace with respect to system B of a bipartite system is defined as

$$\begin{aligned} \text{tr}_B(\rho) &\equiv \sum_{ijkl} c_{ijkl} |a_i\rangle \langle a_j| \otimes \text{tr}(|b_k\rangle \langle b_l|) \\ &= \sum_{ijkl} c_{ijkl} \langle b_l | b_k \rangle |a_i\rangle \langle a_j| \\ &= \sum_{ij} C_{ij} |a_i\rangle \langle a_j| \end{aligned} \quad (2.14)$$

with $C_{ij} = \sum_{kl} c_{ijkl} \langle b_l | b_k \rangle$. The density operator obtained by tracing out one part of the system is called the *reduced density operator*.

We can perform the partial trace on our example state (2.12). The density matrix is

$$\rho = |\alpha|^2 |00\rangle \langle 00| + \alpha\beta^* |00\rangle \langle 11| + \alpha^*\beta |11\rangle \langle 00| + |\beta|^2 |11\rangle \langle 11|, \quad (2.15)$$

so the reduced density matrix becomes

$$\begin{aligned} \rho_A &= \text{tr}_B(\rho) \\ &= |\alpha|^2 \langle 0|0\rangle |0\rangle \langle 0| + \alpha\beta^* \langle 0|1\rangle |0\rangle \langle 1| + \alpha^*\beta \langle 1|0\rangle |1\rangle \langle 0| + |\beta|^2 \langle 1|1\rangle |1\rangle \langle 1| \\ &= |\alpha|^2 |0\rangle \langle 0| + |\beta|^2 |1\rangle \langle 1|, \end{aligned} \quad (2.16)$$

which is just what we expected.

In section 2.2 we saw what we meant by an entangled state as long as the states were pure. But this leaves the question of when a mixed state is entangled. Consider for instance the mixture of two Bell states

$$\rho = \frac{1}{2} |\Phi^+\rangle \langle \Phi^+| + \frac{1}{2} |\Phi^-\rangle \langle \Phi^-|. \quad (2.17)$$

Written out in an orthonormal product basis, this is

$$\begin{aligned} \rho &= \frac{1}{4} (|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|) \\ &\quad + \frac{1}{4} (|00\rangle \langle 00| - |00\rangle \langle 11| - |11\rangle \langle 00| + |11\rangle \langle 11|) \\ &= \frac{1}{2} |00\rangle \langle 00| + \frac{1}{2} |11\rangle \langle 11| \end{aligned} \quad (2.18)$$

which is a mixture of the product states $|00\rangle$ and $|11\rangle$. So the mixed state can be realised by both an ensemble of maximally entangled states and an ensemble of product states. We

say that a mixed state is *separable* if and only if it can be realised as a mixture (convex combination) of locally prepared states⁴. That is, it can be written as⁵

$$\sum_i p_i \rho_i^{(A)} \otimes \rho_i^{(B)} \quad (2.19)$$

where $\{p_i\}$ forms a probability distribution; $p_i \geq 0$ and $\sum_i p_i = 1$. This also means ([HHH96]) that the state can be written as a mixture of pure product states,

$$\sum_{kl} p_{kl} |\psi_k^{(A)}\rangle \langle \psi_k^{(A)}| \otimes |\psi_l^{(B)}\rangle \langle \psi_l^{(B)}|. \quad (2.20)$$

Hence, a mixture of entangled states need not be entangled, but a mixture of separable states is always separable. Mixture is a process which destroys entanglement. This is because by discarding information about which of a number of entangled states the system is in, it can no longer be distinguished from a mixture of separable states.

2.4 The Schmidt decomposition

A useful tool when working with bipartite, pure state entanglement is the *Schmidt decomposition*. It is a decomposition into the biorthogonal basis which gives the smallest possible number of terms for a product basis.

Given a bipartite pure state $|\psi\rangle$, we may write it in terms of some product basis, orthonormal in both subsystems

$$|\psi\rangle = \sum_{i=1}^{d_A} \sum_{j=1}^{d_B} c_{ij} |a_i\rangle \otimes |b_j\rangle. \quad (2.21)$$

The coefficients $\{c_{ij}\}$ may be seen as elements in a $d_A \times d_B$ matrix C . Using the singular value decomposition this may be written as

$$C = UDV \quad (2.22)$$

where D is a $d_A \times d_B$ matrix which is zero except for the diagonal elements which are real and positive, U is a $d_A \times d_A$ unitary matrix and V a $d_B \times d_B$ unitary matrix. This means that c_{ij} may be written as $\sum_{k=1}^{\min(d_A, d_B)} u_{ik} d_{kk} v_{kj}$, so the state vector may be written

$$\begin{aligned} |\psi\rangle &= \sum_{i=1}^{d_A} \sum_{j=1}^{d_B} \sum_{k=1}^{\min(d_A, d_B)} u_{ik} d_{kk} v_{kj} |a_i\rangle \otimes |b_j\rangle \\ &= \sum_{k=1}^{\min(d_A, d_B)} d_{kk} \left(\sum_{i=1}^{d_A} u_{ik} |a_i\rangle \right) \otimes \left(\sum_{j=1}^{d_B} v_{kj} |b_j\rangle \right). \end{aligned}$$

⁴This definition is due to Werner [Wer89] who called them “classically correlated” states.

⁵In fact it was considered sufficient that that such a sum *approximated* the state arbitrarily well. It was shown in [HHH96], though, that in the finite dimensional case any separable state could be expressed in d^2 terms, where d is the dimension of the Hilbert space of the composite system.

We now take as our new basis vectors

$$|a'_k\rangle \equiv \left(\sum_{i=1}^{d_A} u_{ik} |a_i\rangle \right) \quad |b'_k\rangle \equiv \left(\sum_{j=1}^{d_B} v_{kj} |b_j\rangle \right)$$

and define $\lambda_k = d_{kk}$ to obtain the final form

$$|\psi\rangle = \sum_{k=1}^{\min(d_A, d_B)} \lambda_k |a'_k\rangle \otimes |b'_k\rangle \quad (2.23)$$

which is the Schmidt decomposition. What we have achieved is to write any state as a linear combination of maximum $\min(d_A, d_B)$ product vectors. Also, not only are the product vectors in the basis orthogonal, the basis vectors from each subsystem are only used once. These vectors are still orthogonal as we have only performed a unitary transformation on the set of basis vectors, making the set of new basis vectors *biorthogonal*. The coefficients $\{\lambda_k\}$ are called the *Schmidt coefficients*, although in the literature this name is sometimes given to $\{\lambda_k^2\}$.

For many purposes the mere existence of the Schmidt decomposition is enough to make it usable, whereas it sometimes needs to be calculated explicitly. An example of the former is to show that the eigenvalues of the two reduced density matrices of a pure, bipartite state are equal. Actually, it can be seen directly from (2.23) that by forming the density matrix and tracing out *any* one of the two subsystems, the reduced matrix is already on diagonal form with $\{\lambda_k^2\}$ as eigenvalues. It is often easier to calculate the Schmidt coefficients this way than doing the singular value decomposition explicitly.

2.5 Generalised measurements

To facilitate the quantum operations we will generalise the concept of measurements from the way it is usually treated in courses of introductory quantum mechanics. In that formalism, which is called *projective measurements*, the objects representing the measurements are Hermitian operators. The result of the measurement is an eigenvalue of the observable and after the measurement, the new state of the system is in the eigenstate corresponding to the eigenvalue (or projected into the eigenspace if the eigenvalue is degenerate). The probability for an outcome of the measurement is the absolute squared expansion coefficient when the state is expanded in the eigenstates of the observable.

In the general measurement scheme, a measurement is represented by a set of linear operators $\{M_i\}$ called *measurement operators* satisfying the completeness relation

$$\sum_m M_m^\dagger M_m = 1. \quad (2.24)$$

The probability for the outcome to be m is

$$p(m) = \langle \psi | M_m^\dagger M_m | \psi \rangle \quad (2.25)$$

and after the measurement with result m , the new state of the system is

$$\frac{M_m |\psi\rangle}{\sqrt{\langle\psi| M_m^\dagger M_m |\psi\rangle}}. \quad (2.26)$$

We may, at least in theory, implement any measurement that can be described by a set of measurement operators. Thus, we may construct measurements which convert an initial state to a desired state with a certain probability, discarding the system if that particular outcome does not occur. As we will see, this is one of the ideas behind distillation of entanglement.

The projective measurement formalism is a special case of the generalised measurement scheme. The observable, O , is a Hermitian operator and can be decomposed according to the spectral theorem.

$$O = \sum_m \lambda_m |v_m\rangle \langle v_m| \quad (2.27)$$

where $\{|v_m\rangle\}$ are the eigenvectors of the observable, and $\{\lambda_m\}$ the corresponding eigenvalues. By taking as measurement operators $M_m = |v_m\rangle \langle v_m|$, the completeness relation (2.24) is satisfied. The probability for a given outcome, m , of a measurement is

$$\langle\psi| M_m^\dagger M_m |\psi\rangle = \langle\psi| v_m\rangle \langle v_m| v_m\rangle \langle v_m| \psi\rangle = |\langle v_m| \psi\rangle|^2$$

which is exactly the absolute square of the expansion coefficient of $|\psi\rangle$ for that eigenstate. The state after the measurement is

$$\frac{M_m |\psi\rangle}{\sqrt{\langle\psi| M_m^\dagger M_m |\psi\rangle}} = \frac{\langle v_m| \psi\rangle |v_m\rangle}{|\langle v_m| \psi\rangle|} = e^{i\theta} |v_m\rangle$$

with an arbitrary phase θ . Like $|v_m\rangle$, this is an eigenstate of O .

The generalised measurement scheme can easily be further generalised to density operators. For pure states, the generalisation is straightforward. The density operator corresponding to the pure state vector $|\psi\rangle$ is $\rho = |\psi\rangle \langle\psi|$. The probability for a given outcome is of course still the same as in (2.25), but we write it in terms of the density operator:

$$p(m) = \text{tr}(M_m \rho M_m^\dagger) \quad (2.28)$$

When the state changes after a measurement with result m , the density operator corresponding to the state vector (2.26) is obviously

$$\frac{M_m |\psi\rangle \langle\psi| M_m^\dagger}{\langle\psi| M_m^\dagger M_m |\psi\rangle} = \frac{M_m \rho M_m^\dagger}{\text{tr}(M_m \rho M_m^\dagger)} \quad (2.29)$$

The probability (2.28) and new state (2.29) is also valid if the density operator represents a mixed state. To see this, we write the density operator in terms of one of its pure state realisations:

$$\rho = \sum_i p_i |\psi_i\rangle \langle\psi_i| = \sum_i p_i \rho_i \quad (2.30)$$

The joint probability for the system to be in the pure state $|\psi_i\rangle$ and measuring a result m is $p_i p(m|i)$. Summing over all i we get the total probability for getting m as the result of the measurement.

$$\begin{aligned} p(m) &= \sum_i p_i \operatorname{tr}(M_m \rho_i M_m^\dagger) \\ &= \operatorname{tr} \left(M_m \left[\sum_i p_i \rho_i \right] M_m^\dagger \right) \\ &= \operatorname{tr}(M_m \rho M_m^\dagger) \end{aligned}$$

Similarly, the new state after the measurement of the mixed state becomes

$$\rho_{\text{out}} = \sum_i p_i M_m \rho_i M_m^\dagger = M_m \left[\sum_i p_i \rho_i \right] M_m^\dagger = M_m \rho M_m^\dagger.$$

The general measurement scheme covers all operations that can be performed on a quantum system. A unitary transformation is the special case of a one-outcome measurement as the measurement operators then will have to be unitary to satisfy the completeness relation (2.24). Performing a different unitary transformation U_m conditional on the outcome of a measurement described by the operators $\{M_m\}$, is described by the new measurement operators $\{U_m M_m\}$. It is easy to see that these operators satisfy the completeness relation, that the probabilities of the outcomes are the same and that the post measurement state is what we expect, namely $U_m(M_m \rho M_m^\dagger)U_m^\dagger$. Similarly, consecutive measurements can be put together to form a single set of measurement operators, even when the measurements to be done depend on the outcome of the previous ones. When doing this, all possible combinations of measurement outcomes are considered as single outcomes of the total measurement.

In applications it depends on the situation whether we use the full generalised measurement scheme or only consider projective measurements. Projective measurements are sometimes simpler to handle and combined with unitary transformations they describe everything that can be described by generalised measurements. Therefore, both formalisms will be used in the following.

2.6 Transformation of states and LOCC

We will see later that the set of operations that we allow to be performed on a quantum state is essential to the most basic entanglement measures. All operations can be described as generalised measurements. But we will find it useful to describe them as measurements combined with unitary transformations where the next step in a sequence of operations may be decided by a previous measurement outcome.

2.6.1 Local Operations and Classical Communication

In the context of entanglement a subclass of quantum operations is particularly important, namely local quantum operations (LO). When we have a bipartite or multipartite state, the

subsystems may be separated, or at least well isolated from each other. We may therefore only perform operations that act locally on each subsystem. That is, the operators will be of the form $O_A \otimes O_B \otimes \dots$. In addition, the parties sharing a multipartite state are usually allowed to communicate classically, and through this classical communication (CC) they may perform different operations depending on the measurement results of the other parties. This subclass of operations is called *Local Operations and Classical Communication*, usually shortened to *LOCC*. In the literature it is sometimes named otherwise, e.g. QLCC or LQCC. LOCC is important in entanglement theory because entanglement is exactly those correlations that cannot be created using LOCC.

2.6.2 Transformation of a Bell state

Suppose Alice and Bob share a Bell state $|\Phi^+\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$ and want to transform it into the state $\cos\theta|00\rangle + \sin\theta|11\rangle$. One way of doing this is as follows. Alice performs a measurement described by the measurement operators

$$M_1 = \begin{pmatrix} \cos\theta & 0 \\ 0 & \sin\theta \end{pmatrix} \quad M_2 = \begin{pmatrix} \sin\theta & 0 \\ 0 & \cos\theta \end{pmatrix} \quad (2.31)$$

where the local state vectors are denoted as usual:

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad |1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (2.32)$$

The operators clearly satisfies the completeness relation (2.24), as

$$\begin{pmatrix} \cos^2\theta & 0 \\ 0 & \sin^2\theta \end{pmatrix} + \begin{pmatrix} \sin^2\theta & 0 \\ 0 & \cos^2\theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

After the measurement we may have one of two states, depending on the outcome,

$$|\psi_1\rangle = \cos\theta|00\rangle + \sin\theta|11\rangle \quad |\psi_2\rangle = \sin\theta|00\rangle + \cos\theta|11\rangle$$

where $|\psi_1\rangle$ is the desired output state. If the result of the measurement is 2, leaving the system in the state $|\psi_2\rangle$, Alice applies the X operator (also called bit flip operator) to her system converting $|0\rangle$ into $|1\rangle$ and vice versa. Then she tells Bob to do the same through the classical communication channel. The resulting state is $|\psi_1\rangle$ and their common goal is achieved.

The transformation we just did, transforms a maximally entangled state into a less or equally entangled state. For $\theta = 0$ and $\theta = \pi/2$ the output state is a product state, which is not entangled at all, whereas for $\theta = \pi/4$ and $\theta = 3\pi/4$ the output state is also a Bell state.

2.6.3 Concentrating partial pure state entanglement

We may also try to do the transformation the other way around. We begin with the state $|\psi_1\rangle$, where we for simplicity suppose that $0 < \theta < \pi/4$, and want to end up with the Bell state $|\Phi^+\rangle$. The lower boundary of this interval is a product state for $\theta = 0$ which cannot

possibly be turned into an entangled state by LOCC. The upper boundary $\theta = \pi/4$ is the desired output state and will need no transformation.

The problem is now to construct local measurement operators which give $|\Phi^+\rangle$ as a post measurement state. The operator

$$M_1 = \text{const} \cdot \begin{pmatrix} \frac{1}{\cos \theta} & 0 \\ 0 & \frac{1}{\sin \theta} \end{pmatrix}$$

will give the desired output, and we want the constant to be as large as possible to maximise the probability for this output. Still, for the completeness relation (2.24) to be satisfied none of the diagonal elements can be greater than 1. This is because for any M_m , $M_m^\dagger M_m$ has nonnegative diagonal elements, and each diagonal element needs to sum up to 1 to get the identity operator in (2.24). In the valid range of θ , $\cos \theta > \sin \theta$, so the maximal constant is $\sin \theta$. This gives the optimal measurement operator for the successful outcome

$$M_1 = \begin{pmatrix} \tan \theta & 0 \\ 0 & 1 \end{pmatrix} = \tan \theta |0\rangle \langle 0| + |1\rangle \langle 1|.$$

The remaining set of matrices representing the measurement operators need to have a vanishing element to the lower right, and the remaining elements need to be such that the completeness relation is satisfied. The easiest way to do this is to simply add one more operator, namely

$$M_2 = \begin{pmatrix} \sqrt{1 - \tan^2 \theta} & 0 \\ 0 & 0 \end{pmatrix} = \sqrt{1 - \tan^2 \theta} |0\rangle \langle 0|.$$

The new states after the measurement are proportional to

$$\begin{aligned} M_1 |\psi_1\rangle &= \sin \theta |00\rangle + \sin \theta |11\rangle \\ M_2 |\psi_1\rangle &= \cos \theta \sqrt{1 - \tan^2 \theta} |00\rangle. \end{aligned}$$

The probability for the two outcomes are

$$\begin{aligned} p(1) &= \langle \psi_1 | M_1^\dagger M_1 | \psi_1 \rangle = \sin^2 (\langle 00 | 00 \rangle + \langle 11 | 11 \rangle) = 2 \sin^2 \theta \\ p(2) &= \langle \psi_1 | M_2^\dagger M_2 | \psi_1 \rangle = \cos^2 \theta (1 - \tan^2 \theta) = \cos^2 \theta - \sin^2 \theta = 1 - 2 \sin^2 \theta \end{aligned}$$

and the normalised new states are

$$\begin{aligned} |\psi_{\text{out},1}\rangle &= \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle) = |\Phi^+\rangle \\ |\psi_{\text{out},2}\rangle &= |00\rangle. \end{aligned}$$

We see that there is a certain probability for converting a partially entangled state into a maximally entangled state, but the less entangled the initial state is, the smaller is the probability. This is consistent with the fact that LOCC cannot increase the expected entanglement. Still, we may perform a measurement which may give us a more entangled state at the risk of losing the entanglement we had. If we have a large number of bipartite two-qubit systems, we may apply the above procedure and discard the systems where the output is a product state, thereby effectively *concentrating* entanglement.

2.6.4 Distilling entanglement from mixed states

Also mixed state entanglement may be concentrated, or *distilled*. This will be important for practical applications, as the channels used to create entanglement in general will be noisy. A collection of distillation protocols was introduced in [BBP⁺96] and [BDSW96]. We will take a closer look at a slightly modified version of a distillation protocol introduced in the latter, which for certain mixed states can yield pure Bell states as output.

Consider a mixed state, as always shared between Alice and Bob, which can be represented as a collection of $|\Phi^+\rangle$ states contaminated with a portion of $|01\rangle$ states. The corresponding density operator is then

$$\begin{aligned}\rho &= (1-p) |01\rangle \langle 01| + p |\Phi^+\rangle \langle \Phi^+| \\ &= (1-p) |01\rangle \langle 01| + \frac{p}{2} |00\rangle \langle 00| + \frac{p}{2} |00\rangle \langle 11| + \frac{p}{2} |11\rangle \langle 00| + \frac{p}{2} |11\rangle \langle 11|.\end{aligned}$$

Alice and Bob now pick two pairs from the ensemble. Each of them will perform a joint operation on the two subsystems on their side. Then they measure one of the pairs, thereby destroying any entanglement in that pair, to decide if the other pair is in the desired output state.

The composite system of the two pairs is described by the density operator

$$\begin{aligned}\rho^{\otimes 2} &= \\ (1-p)^2 &|0101\rangle \langle 0101| + \frac{p(1-p)}{2} |0100\rangle \langle 0100| + \frac{p(1-p)}{2} |0100\rangle \langle 0111| + \frac{p(1-p)}{2} |0111\rangle \langle 0100| + \frac{p(1-p)}{2} |0111\rangle \langle 0111| \\ &+ \frac{p(1-p)}{2} |0001\rangle \langle 0001| + \frac{p^2}{4} |0000\rangle \langle 0000| + \frac{p^2}{4} |0000\rangle \langle 0011| + \frac{p^2}{4} |0011\rangle \langle 0000| + \frac{p^2}{4} |0011\rangle \langle 0011| \\ &+ \frac{p(1-p)}{2} |0001\rangle \langle 1101| + \frac{p^2}{4} |0000\rangle \langle 1100| + \frac{p^2}{4} |0000\rangle \langle 1111| + \frac{p^2}{4} |0011\rangle \langle 1100| + \frac{p^2}{4} |0011\rangle \langle 1111| \\ &+ \frac{p(1-p)}{2} |1101\rangle \langle 0001| + \frac{p^2}{4} |1100\rangle \langle 0000| + \frac{p^2}{4} |1100\rangle \langle 0011| + \frac{p^2}{4} |1111\rangle \langle 0000| + \frac{p^2}{4} |1111\rangle \langle 0011| \\ &+ \frac{p(1-p)}{2} |1101\rangle \langle 1101| + \frac{p^2}{4} |1100\rangle \langle 1100| + \frac{p^2}{4} |1100\rangle \langle 1111| + \frac{p^2}{4} |1111\rangle \langle 1100| + \frac{p^2}{4} |1111\rangle \langle 1111|\end{aligned}$$

where the first and third qubit are in Alice's possession and the second and fourth are in Bob's possession. The two first qubits are the first pair, and the two last are the second pair. It may be written symbolically as $|A_1 B_1 A_2 B_2\rangle$. Both Alice and Bob applies a CNOT gate to the two qubits in their possession, using the qubit from the first pair as a control bit and the second one as the target. A CNOT gate is a unitary transformation which does nothing when the control bit is zero and flips the target bit if the control bit is set to 1. The operator is written

$$\begin{aligned}U_{\text{CNOT}} &= |00\rangle \langle 00| + |01\rangle \langle 01| + |10\rangle \langle 11| + |11\rangle \langle 10| \\ &= |0\rangle \langle 0| \otimes \mathbb{I} + |1\rangle \langle 1| \otimes X\end{aligned}$$

where X is the usual bit flip operator, or Pauli X -operator. But as Alice's operator works on the first and third qubit while Bob's operator works on the second and fourth, in our basis the operators are

$$\begin{aligned}U_{\text{CNOT}}(1,3) &= |0\rangle \langle 0| \otimes \mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} + |1\rangle \langle 1| \otimes \mathbb{I} \otimes X \otimes \mathbb{I} \\ U_{\text{CNOT}}(2,4) &= \mathbb{I} \otimes |0\rangle \langle 0| \otimes \mathbb{I} \otimes \mathbb{I} + \mathbb{I} \otimes |1\rangle \langle 1| \otimes \mathbb{I} \otimes X\end{aligned}$$

Operationally it only amounts to flipping the third qubit of $\rho^{\otimes 2}$ in all terms where the first qubit is 1 and flipping the fourth qubit where the second qubit is 1. This leaves the new state

$$\begin{aligned} \rho_{\text{out}} = & U_{\text{CNOT}}(1, 3)U_{\text{CNOT}}(2, 4)\rho^{\otimes 2}U_{\text{CNOT}}^\dagger(2, 4)U_{\text{CNOT}}^\dagger(1, 3) = \\ & (1-p)^2 |0100\rangle\langle 0100| + \frac{p(1-p)}{2} |0101\rangle\langle 0101| + \frac{p(1-p)}{2} |0110\rangle\langle 0110| + \frac{p(1-p)}{2} |0110\rangle\langle 0101| + \frac{p(1-p)}{2} |0110\rangle\langle 0110| \\ & + \frac{p(1-p)}{2} |0001\rangle\langle 0001| + \frac{p^2}{4} |0000\rangle\langle 0000| + \frac{p^2}{4} |0000\rangle\langle 0011| + \frac{p^2}{4} |0011\rangle\langle 0000| + \boxed{\frac{p^2}{4} |0011\rangle\langle 0011|} \\ & + \frac{p(1-p)}{2} |0001\rangle\langle 1110| + \frac{p^2}{4} |0000\rangle\langle 1111| + \frac{p^2}{4} |0000\rangle\langle 1100| + \boxed{\frac{p^2}{4} |0011\rangle\langle 1111|} + \frac{p^2}{4} |0011\rangle\langle 1100| \\ & + \frac{p(1-p)}{2} |1110\rangle\langle 0001| + \frac{p^2}{4} |1111\rangle\langle 0000| + \boxed{\frac{p^2}{4} |1111\rangle\langle 0011|} + \frac{p^2}{4} |1100\rangle\langle 0000| + \frac{p^2}{4} |1100\rangle\langle 0011| \\ & + \frac{p(1-p)}{2} |1110\rangle\langle 1110| + \boxed{\frac{p^2}{4} |1111\rangle\langle 1111|} + \frac{p^2}{4} |1111\rangle\langle 1100| + \frac{p^2}{4} |1100\rangle\langle 1111| + \frac{p^2}{4} |1100\rangle\langle 1100|. \end{aligned}$$

In this state Alice and Bob both measure the state of their qubits belonging to the target pair. This can be done formally with the set of measurement operators

$$M_{00} = \mathbb{I} \otimes \mathbb{I} \otimes |0\rangle\langle 0| \otimes |0\rangle\langle 0| \quad (2.33a)$$

$$M_{01} = \mathbb{I} \otimes \mathbb{I} \otimes |0\rangle\langle 0| \otimes |1\rangle\langle 1| \quad (2.33b)$$

$$M_{10} = \mathbb{I} \otimes \mathbb{I} \otimes |1\rangle\langle 1| \otimes |0\rangle\langle 0| \quad (2.33c)$$

$$M_{11} = \mathbb{I} \otimes \mathbb{I} \otimes |1\rangle\langle 1| \otimes |1\rangle\langle 1|, \quad (2.33d)$$

and is equivalent to measuring the observable $\mathbb{I} \otimes \mathbb{I} \otimes Z_3 \otimes Z_4$ as the measurement operators are constructed from eigenvectors of this observable. The interesting outcome is when both qubits are measured to 1. The probability for that is

$$p(11) = \text{tr} \left(M_{11} \rho_{\text{out}} M_{11}^\dagger \right) \quad (2.34)$$

where only the boxed terms of ρ_{out} survive being operated on by the M_{11} and of them only the two diagonal ones give a nonzero contribution when the trace is taken, giving a probability of $\frac{p^2}{2}$. The new state after the measurement is

$$\begin{aligned} \frac{M_{11} \rho_{\text{out}} M_{11}^\dagger}{p(11)} &= \frac{1}{2} \left(|00\rangle\langle 00| + |00\rangle\langle 11| + |11\rangle\langle 00| + |11\rangle\langle 11| \right) \otimes |11\rangle\langle 11| \\ &= |\Phi^+\rangle\langle \Phi^+| \otimes |11\rangle\langle 11|. \end{aligned} \quad (2.35)$$

So by using two entangled pairs in a mixed state Alice and Bob have obtained a pure state where one of the pairs is in a Bell state. The probability for this to happen is $\frac{p^2}{2}$, so on average we get a yield of $\frac{p^2}{4}$ Bell states per input pair.

Some additional Bell states may be squeezed out of the mixed states after using the above protocol. The measurement result 00, with a probability $\frac{3}{2}p^2 - 2p + 1$, leaves the pair of control qubits in the state

$$\rho_{00} = \frac{2(1-p)^2}{3p^2 - 4p + 2} |01\rangle\langle 01| + \frac{p^2}{3p^2 - 4p + 2} |\Phi^+\rangle\langle \Phi^+| \quad (2.36)$$

which is on the same form as the original input state with a new mixing ratio

$$p' = \frac{p^2}{3p^2 - 4p + 2}. \quad (2.37)$$

These pairs may be run through the same procedure once more with a yield of $\frac{p'^2}{4}$, increasing the total yield with an amount of $\frac{1}{2}p(00) \cdot \frac{p'^2}{4}$, where the factor $\frac{1}{2}$ comes from the fact that we use two pairs and only get one out. This procedure may be iterated, and the total yield converges quickly so only few iterations are necessary in practice. Figure 2.2 shows the total yield calculated numerically as a function of p .




Figure 2.2: The yield of the distillation protocol for 1, 2 and 3 iterations. The dotted line is the asymptotic limit.

In general, distillation protocols do not yield pure state pairs after a finite number of iterations. Likewise, in the above protocol this only happens when the state can be written as a mixture of a Bell state and another special state. A general feature, though, is the application of the bilateral CNOT unitary transformation on two pairs.

The purity of the output states is measured by the *fidelity* of the state ρ , defined as⁶

$$F \equiv \langle \Phi^+ | \rho | \Phi^+ \rangle \quad (2.38)$$

where $|\Phi^+\rangle$ is the desired output state. Applying a distillation protocol to an ensemble of states, leaves a subensemble where the fidelity is higher, along with a subensemble where the fidelity is lower, often zero. In the limit of reapplying the procedure to the higher-fidelity states an infinite number of times, the fidelity approaches unity. This comes at a cost,

⁶Sometimes the fidelity is defined as the square root of this.

however, as the yield decreases as we demand higher fidelity. For the simplest protocols the yield approaches zero as the fidelity approaches unity. But by combining protocols and using the already distilled states as a resource, the limit can be made finite.

Chapter 3

Characterisation of bipartite entanglement

Before moving on to quantify the entanglement in a bipartite state, we will try to characterise it more qualitatively. There are a number of questions which does not have an immediate answer. Given a state, how can we tell wheter it is entangled? Is there more than one type of entanglement? Can all entangled states be distilled to Bell states?

3.1 Pure states

When we are given a bipartite state the first question of interest is whether it is entangled or not. This is the question of *separability*. A separable state is a state which is not entangled, so each of the subsystems can be given a description on its own. When it comes to pure states, a state is separable if and only if it is a product state. As indicated in 2.2, a pure state is a product state if it can be written as

$$|\Psi_{\text{prod}}\rangle = |\psi\rangle_A \otimes |\phi\rangle_B \quad (3.1)$$

for any pure states $|\psi\rangle_A$ and $|\phi\rangle_B$ belonging to Alice and Bob, respectively. If the state is not a product state, it is an entangled state.

In practice, though, one needs to find the right basis vectors to be able to write a product state as in (3.1). A simpler criterion is desired. The Schmidt decomposition (section 2.4) is helpful in this respect. The number of terms in the Schmidt decomposition is called the *Schmidt number*, sometimes also *Schmidt rank*. That is, the number n in

$$|\psi\rangle = \sum_{k=1}^n \lambda_k |a'_k\rangle \otimes |b'_k\rangle. \quad (3.2)$$

The Schmidt number is at most $\min(d_A, d_B)$, where d_A (d_B) is the dimension of subsystem A (B). But when some of the Schmidt coefficients are zero, those terms may be skipped giving a lower Schmidt number. Remember that the Schmidt coefficients are simply the square root of the eigenvalues of any of the reduced density matrices. Hence, the Schmidt number is simply the number of nonzero eigenvalues of ρ_A or ρ_B .

What the Schmidt number tells us physically is basically how many degrees of freedom that are entangled. And as a product state can be written as (3.1) it is obvious that its Schmidt number is 1. For pure states this is a necessary and sufficient criterion for separability.

It turns out that entanglement from any pure entangled state can be concentrated into Bell states. This may be done by using the procedure similar to the one described in section 2.6.3, or a more advanced procedure that operates on more partially entangled pairs at the same time to give a higher yield (see e.g. [BBPS96]). Bell states may again be used to produce other entangled states by means of LOCC. Because of this, and the easy identification through the Schmidt number, the only classification we make for bipartite pure states is in separable and entangled states.

3.2 Mixed states

Some properties of mixed states were discussed in section 2.3. The structure of mixed state bipartite entanglement is richer than its pure state equivalent. For instance there are entangled states that cannot be distilled to Bell states, and cannot be put to use in any of the applications where entanglement is a vital resource. Also, the more difficult task of finding out whether a given mixed state is separable or not gives rise to classes of states which can be identified by other criteria.

3.2.1 Separable states

Recall from section 2.3 that a bipartite state is called separable if it can be realised as a mixture of product states,

$$\sum_i p_i \rho_i^{(A)} \otimes \rho_i^{(B)} \quad (3.3)$$

with $p_i \geq 0$ and $\sum_i p_i = 1$. Given a density matrix in some basis the task of finding local density matrices to satisfy (3.3) can be enormous. Even for the simplest system which may be entangled, a pair of qubits, a density operator would typically look like

$$\begin{aligned} \rho = & \frac{3}{24} |00\rangle \langle 00| + \frac{\sqrt{2}}{24} i |00\rangle \langle 01| + \frac{\sqrt{2}}{12} i |00\rangle \langle 11| \\ & - \frac{\sqrt{2}}{24} i |01\rangle \langle 00| + \frac{1}{4} |01\rangle \langle 01| - \frac{\sqrt{2}}{12} i |01\rangle \langle 10| \\ & + \frac{\sqrt{2}}{12} i |10\rangle \langle 01| + \frac{5}{24} |10\rangle \langle 10| - \frac{\sqrt{2}}{24} i |10\rangle \langle 11| \\ & - \frac{\sqrt{2}}{12} i |11\rangle \langle 00| + \frac{\sqrt{2}}{24} i |11\rangle \langle 10| + \frac{5}{12} |11\rangle \langle 11| \end{aligned} \quad (3.4)$$

or in matrix representation in the $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ basis,

$$\rho = \begin{pmatrix} \frac{3}{24} & \frac{\sqrt{2}}{24} i & 0 & \frac{\sqrt{2}}{12} i \\ -\frac{\sqrt{2}}{24} i & \frac{1}{4} & -\frac{\sqrt{2}}{12} i & 0 \\ 0 & \frac{\sqrt{2}}{12} i & \frac{5}{24} & -\frac{\sqrt{2}}{24} i \\ -\frac{\sqrt{2}}{12} i & 0 & \frac{\sqrt{2}}{24} i & \frac{5}{12} \end{pmatrix}. \quad (3.5)$$

It is not easy to see that this can be written as (3.3). Still, it is constructed as a mixture of two product states. It can be written as

$$\begin{aligned}\rho &= p_1 \rho_1^{(A)} \otimes \rho_1^{(B)} + p_2 \rho_2^{(A)} \otimes \rho_2^{(B)} \\ &= \frac{1}{2} \left(\frac{1}{2} |1\rangle \langle 1| + \frac{1}{2} |- \rangle \langle -| \right) \otimes |\psi_{\frac{2}{3}}^+\rangle \langle \psi_{\frac{2}{3}}^+| + \frac{1}{2} \left(\frac{1}{2} \mathbb{I}/2 + \frac{1}{2} |+\rangle \langle +| \right) \otimes |\psi_{\frac{2}{3}}^-\rangle \langle \psi_{\frac{2}{3}}^-| \quad (3.6)\end{aligned}$$

where

$$|\psi_{\frac{2}{3}}^\pm\rangle = \frac{1}{\sqrt{3}} |0\rangle \pm \sqrt{\frac{2}{3}} i |1\rangle$$

It could of course be realised by a variety of other mixtures as well.

This example shows that we could well use an operational criterion for separability, like the Schmidt number for pure states. In general, no easily computable necessary and sufficient criterion is known. Still, we have a necessary condition which is easy to compute. This is known as the *PPT¹-criterion* or *Peres-Horodecki criterion* [Per96, HHH96]. It states that if the state is separable, then the partial transpose of the density operator with respect to one subsystem is positive. A transpose of an operator needs to be taken with respect to a basis, and the resulting operator depends on the basis. Actually, if A^T is the transpose of the operator with respect to one basis, any operator of the form $UA^TU^\dagger$ for unitary U satisfying $U = U^T$ (in the same basis) is a transpose of the operator with respect to another basis. However, since the different bases are connected by a unitary transformation, the eigenvalues are independent of the basis. Therefore, any orthonormal basis will do for taking the transposition. More formally, if we choose an orthonormal product basis $\{|v_i v_j\rangle\} \equiv \{|v_i\rangle \otimes |v_j\rangle\}$ for the state ρ , the partial transpose ρ^{T_B} is defined by its matrix elements

$$\rho_{m\mu, n\nu}^{T_B} = \langle v_m v_\mu | \rho^{T_B} | v_n v_\nu \rangle = \rho_{m\nu, n\mu}. \quad (3.7)$$

In the example matrix (3.5) taking the partial transpose of system B corresponds to transposing within each of the four 2×2 blocks, whereas switching the upper right block with the lower left block corresponds to transposing system A .

For systems of dimension 2×2 and 2×3 the PPT-criterion is also sufficient for the state to be separable [HHH96]. For larger systems, though, there exist entangled PPT states [Hor97]. The PPT-criterion is still so useful that the class of PPT states is an important class of states, strictly larger than the class of separable states. Anything that can be proved for all PPT states, is automatically true for separable states.

3.2.2 Free and bound entanglement

Unlike pure states, not all entangled mixed states can be distilled into Bell states [HHH98]. This is an important distinction, because for many applications entanglement must be distilled in order to be useful. Entanglement that can be distilled is called *free entanglement* and entanglement that cannot be distilled is called *bound entanglement*. Bound entangled states cannot be written as a convex combination of product states (3.3) and thus cannot be prepared locally by two classically communicating parties, but its entanglement is well

¹Positive Partial Transpose

hidden and difficult to detect. It has been shown that for a pair of qubits all entangled states can be distilled [HHH97].

A connection between free entangled states and PPT states was found in [HHH98]. It says that no free entangled state is PPT. Hence, all PPT entangled states are bound entangled states. This result was actually the first evidence of the existence of bound entangled states.

This raises the question of whether a nonpositive partial transpose (NPPT) state is always distillable. The answer to this question is not known, but there are indications that NPPT bound entangled states do exist [DSS⁺00, DCLB00].

It has been proven that bound entanglement is useless for quantum teleportation [HHH99a], and it can be proved for other applications as well. One has found, though, that bound entanglement is not completely useless after all. For certain tasks it can be activated [HHH99b] to facilitate distillation of other quantum states. It has even been shown that a small amount of PPT bound entanglement as a resource makes it possible to distill entanglement from any NPPT state [VW02b]. This means that if NPPT bound entangled states really exist, entanglement can be distilled by putting together two states that are not distillable by themselves.

Chapter 4

Measures of bipartite entanglement

Entanglement can be utilised for performing many tasks which are impossible without it, or enhance the performance of other tasks. In most of these tasks entanglement is consumed, so we trade entanglement for something else. It is clear from this that we may treat entanglement as a *resource*, just as energy is a resource needed for certain tasks. This is the reason why we want to *quantify* the entanglement in an entangled state. Given a state and a task that consumes entanglement, how much can we achieve? How well can we do? It is not obvious that an entangled state that performs a given task better than another is will be the better resource for another task. Indeed, this is only true in a very limited context. Because of this, we have in general many ways to quantify the nonlocal resources (or entanglement) in a quantum state.

An entanglement measure $E(\cdot)$ is a functional that takes a quantum state of a multipartite system to a nonnegative real number. In this chapter we only consider bipartite systems, hence

$$E : \mathcal{D}(\mathcal{H}) \rightarrow \mathbb{R}^+ \quad (4.1)$$

where $\mathcal{D}(\mathcal{H})$ is the set of density operators on the Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$. There are in principle two ways to quantify the entanglement in a state. *Operational measures* are based on how well a certain task can be performed, usually compared to Bell states. The other way, which gives rise to *abstract measures*, is to work from a set of natural axioms we believe an entanglement measure should satisfy, and look for functionals that satisfy the axioms.

There are also two regimes to consider. In the *finite regime* we consider the resources contained in a single quantum system, whereas in the *asymptotic regime* we take into account an infinite number of systems in the same state and consider the the resources *per system*.

4.1 Pure states

Before treating the general case of mixed states we will consider the simpler case of entanglement measures of pure states. The treatment is facilitated by the fact that a pure state contains no classical correlations between the subsystems, so any correlation present must be of quantum nature. As mentioned in section 2.3, even if an entangled state is

pure the states of the subsystems – described locally by the reduced density operator – are mixed. For pure states, then, the amount of “mixedness” turn out to be a good measure of entanglement.

4.1.1 von Neumann entropy

Our measure for how mixed a quantum state ρ is, will be the *von Neumann entropy*, defined as

$$S(\rho) \equiv -\text{tr}(\rho \log \rho) \quad (4.2)$$

where we take the logarithm base-2 as is the custom in information theory. It is most easily calculated from the nonzero eigenvalues λ_i of ρ as

$$S(\rho) = -\sum_i \lambda_i \log \lambda_i. \quad (4.3)$$

The von Neumann entropy is often seen as a generalisation of the Shannon entropy from classical information theory. Although it is true that Shannon entropy arises as a special case in quantum information theory when only orthogonal states are considered, the historical influence is the other way around. von Neumann introduced his entropy in 1932 [vN32] and Shannon published his mathematical theory of communication in 1948 [Sha48]. In fact, it was von Neumann who suggested that Shannon should name his uncertainty function “entropy” as that was already used in statistical physics [TM71, p. 180].

What the von Neumann entropy essentially describes, is the uncertainty in a quantum state. It is zero for pure states, and smaller for a mixture of two nonorthogonal states than a similar mixture of two orthogonal states. It is not only an analogue to the entropy known from thermodynamics, it is the same quantity (up to a constant with the definition (4.2)). Indeed, it was derived using notions such as the ideal gas law and other laws of thermodynamics [vN32, p. 191-202]¹.

4.1.2 Reduced von Neumann entropy – entropy of entanglement

For the purpose of quantifying entanglement, the important quantity is the *reduced* von Neumann entropy. This is the von Neumann entropy of the reduced density matrix. We saw an example in section (2.3) that the reduced density operator of a pure entangled state represents a mixed state. The von Neumann entropy of *either* of the subsystems (the entropies are equal) is a good measure of entanglement. Because of this, the reduced von Neumann entropy is also known under the name *entropy of entanglement* E_E .

It is easy to see that the reduced von Neumann entropy is equal for both reduced density matrices. The von Neumann entropy only depends on the eigenvalues of the density matrix (4.3) and because of the existence of the Schmidt decomposition, the eigenvalues are equal, as was mentioned in section 2.4.

We haven’t yet pointed out what constitutes a good measure, but the entropy of entanglement has some properties we find natural. (i) It is zero for any product state, (ii) it is maximal when the reduced density matrix is completely mixed, e.g. when the subsystems

¹Pages 359-379 in the far worse typeset English translation [vN55].

have no individual properties (for the degrees of freedom considered) and (iii) it is invariant under local unitary transformations. The above requirements were introduced in [SM95], but requirements have later been refined and placed on firmer ground for both pure and mixed states.

The entropy of entanglement was first introduced as a measure of entanglement in [BBP⁺96]. It is an abstract measure in the sense that it satisfies some requirements, and it does not have an immediate operational interpretation. We will now take a look at two pure state entanglement measures which have a direct operational meaning.

4.1.3 Entanglement cost and distillable entanglement

We saw in section 2.6.3 an example of a concentration of pure state entanglement. We turned a partly entangled two-qubit state into a Bell state with a yield depending on how entangled the initial state was. There are other protocols which give a higher yield if we have more than one pair to operate on (see e.g. [BBPS96]). The *distillable entanglement* E_d is defined as the maximum yield of Bell states that can be obtained, optimised over all possible LOCC protocols. The distillable entanglement is also sometimes called *entanglement of distillation*.

The *entanglement cost* E_c is the dual to the distillable entanglement. Two separated parties cannot prepare an entangled state if they can only communicate classically. But if they have some entangled state in some standard form like Bell states, they can convert those into the desired entangled state. Thus, the entanglement cost is defined as the minimum number of Bell states needed to create a given state by means of LOCC.

The above definitions were implicitly given for the finite regime. It turns out that in this regime E_d and E_c are hard to calculate even though the states are pure. What is clear intuitively is that for any state

$$E_d \leq E_c. \quad (4.4)$$

If not, one could create entanglement by means of LOCC by converting Bell states to a state not satisfying (4.4) and converting them back again.

In the asymptotic limit, the results are known. We define the regularised (or asymptotic) versions of distillable entanglement and entanglement cost as

$$E_D(|\psi\rangle) \equiv E_d^\infty(|\psi\rangle) \equiv \lim_{n \rightarrow \infty} \frac{E_d(|\psi\rangle^{\otimes n})}{n}; \quad (4.5a)$$

$$E_C(|\psi\rangle) \equiv E_c^\infty(|\psi\rangle) \equiv \lim_{n \rightarrow \infty} \frac{E_c(|\psi\rangle^{\otimes n})}{n}. \quad (4.5b)$$

In this case, it was shown in [BBPS96] that both the distillable entanglement and entanglement cost is equal to the entropy of entanglement, $E_D(|\psi\rangle) = E_C(|\psi\rangle) = E_E(|\psi\rangle)$. In fact, if two states $|\psi_1\rangle$ and $|\psi_2\rangle$ have the same entropy of entanglement, they can be interconverted with efficiency approaching unity as $n \rightarrow \infty$. If they do not have the same entropy of entanglement, $|\psi_1\rangle$ can be converted into $|\psi_2\rangle$ with asymptotic yield $E_E(|\psi_1\rangle)/E_E(|\psi_2\rangle)$. It has even been shown that this conversion can be done such that the amount of classical communication required per Bell state produced approaches zero [LP99] (it scales as $\sqrt{n}$).

Despite the fact that the entropy of entanglement did not have an operational interpretation *ab initio*, we see that it indeed plays a fundamental role. The fact that the entropy

of entanglement can be asymptotically conserved in the conversion of states using LOCC, places huge constraints on other candidates for pure state asymptotic entanglement measures. A measure would not live up to our expectations if we could increase it using only LOCC. If another measure were to give two different values for states with the same entropy of entanglement, it would be possible to increase it by LOCC by converting the state with lower value to the one with higher value. The conversion between states with conversion rate $E_E(|\psi_1\rangle)/E_E(|\psi_2\rangle)$ imposes even stronger constraints, so we would expect any asymptotic pure state measure to coincide with the entropy of entanglement. The exact conditions for this to happen are discussed in the next section.

4.1.4 The uniqueness theorem for measures of entanglement

The fact that the pure state asymptotic entanglement cost and distillable entanglement coincide with the entropy of entanglement is a special case of a more general property of entanglement measures. The *uniqueness theorem for entanglement measures* states that any pure state measure of entanglement that satisfies certain natural criteria coincides with the entropy of entanglement. The criteria are considered to be too strict for the finite regime, but all natural asymptotic measures satisfy the conditions.

The uniqueness theorem has been developed gradually in the literature, in the beginning only as an intuitive idea of the same character as given in the previous section [PR97]. Later, uniqueness was proved from a set of conditions that measures should satisfy [Vid00, HHH00]. In [DHR02] the minimal conditions for the theorem to hold were found. The conditions come in different versions, some weaker than others. It turns out that some of the stronger conditions are not necessary for the theorem, but are in fact implied by the weaker conditions. The strongest versions are listed first.

(P0) If $|\psi\rangle$ is separable (i.e. a product state), then $E(|\psi\rangle) = 0$

(P1a) (Normalisation) For the maximally entangled state in $d \times d$ dimensions

$$|\Phi_d^+\rangle = \sum_{i=1}^d \frac{1}{\sqrt{d}} |i\rangle_A \otimes |i\rangle_B \quad (4.6)$$

where the two sets $\{|i\rangle_A\}$ and $\{|i\rangle_B\}$ are orthonormal bases for the two subsystems, $E(|\Phi_d^+\rangle) = \log d$, where we as usual take the logarithm base-2.

(P1b) For the Bell state $|\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, $E(|\Phi^+\rangle) = 1$. The previous condition is simply the generalisation of this.

(P2) For any operation Λ that can be implemented by means of LOCC and any $|\psi\rangle$ such that $\Lambda(|\psi\rangle)$ is a pure state, $E(\Lambda(|\psi\rangle)) \leq E(|\psi\rangle)$.

(P3) (Continuity) Let $\{|\psi_n\rangle\}$ and $\{|\phi_n\rangle\}$ be sequences of pure bipartite states living on the sequence of Hilbert spaces $\{\mathcal{H}_n\}$. For all such sequences such that $\| |\psi_n\rangle \langle \psi_n| - |\phi_n\rangle \langle \phi_n| \|_1 \rightarrow 0$ where $\|\cdot\|_1$ is the trace norm $\|A\|_1 = \text{tr}(\sqrt{A^\dagger A})$,

$$\frac{E(|\psi\rangle) - E(|\phi\rangle)}{1 + \log(\dim \mathcal{H}_n)} \rightarrow 0. \quad (4.7)$$

(P4a) (Weak additivity) For all pure states $|\psi\rangle$ and $n \geq 1$,

$$\frac{E(|\psi\rangle^{\otimes n})}{n} = E(|\psi\rangle) \quad (4.8)$$

(P4b) (Asymptotic weak additivity) Given $\epsilon > 0$ and a pure state $|\psi\rangle$, there exists an integer $N > 0$ such that for all $n \geq N$,

$$\frac{E(|\psi\rangle^{\otimes n})}{n} - \epsilon \leq E(|\psi\rangle) \leq \frac{E(|\psi\rangle^{\otimes n})}{n} + \epsilon. \quad (4.9)$$

(P4c) (Existence of a regularisation) For all bipartite pure states $|\psi\rangle$, the limit

$$E^\infty(|\psi\rangle) \equiv \lim_{n \rightarrow \infty} \frac{E(|\psi\rangle^{\otimes n})}{n} \quad (4.10)$$

exists.

With these conditions given, the uniqueness theorem for entanglement measures states [DHR02] that for a functional E on pure states, the following are equivalent:

- (i) E satisfies (P1b), (P2), (P3) and (P4b).
- (ii) E satisfies (P0), (P1a), (P2), (P3) and (P4a).
- (iii) E coincides with the entropy of entanglement $E = E_E$.

On the other hand, if E satisfies (P0), (P1a), (P2) and (P3), it automatically satisfies (P4c), and $E^\infty = E_E$.

(i) represents the weakest set of conditions. When they are satisfied, the stronger conditions in (ii) are also satisfied. The connection is that (P4a) and (P4b) turn out to be equivalent ((P4a) $\Rightarrow$ (P4b) is obvious) and (P1b), (P2) and (P4a) together implies (P0) and (P1a). The uniqueness for the asymptotic regime, previously mentioned, is concretised by the last sentence.

4.2 Mixed states

The measures of pure state bipartite entanglement constitute a very limited subset of the measures of generally mixed states. For mixed states there is no unique way to quantify the nonlocal quantum resources even in the asymptotic regime. In the pure state asymptotic limit, the entropy of entanglement imposed a total order on the set of states. Any state could be converted into another state with equal or less entanglement. In the mixed state case there are states where neither can be converted into the other, giving only a partial order on the set of states. (For a longer discussion on entanglement measures as ordering of the states, see [MSV04].) This makes the structure of mixed state entanglement richer than that of pure states.

4.2.1 Entanglement cost and distillable entanglement for mixed states

The entanglement cost and distillable entanglement can easily be defined for mixed states in the same way as for pure states. The entanglement cost E_c is the minimum number of Bell states needed to produce the state by means of LOCC and the distillable entanglement E_d

is the maximum number of Bell states that can be distilled by an optimal LOCC distillation protocol. However, a perfect conversion between mixed states in the finite regime is usually not possible, so in the following we will only consider the asymptotic versions E_D and E_C . There is a tweak, though, compared to the pure state case. As mentioned in the discussion of entanglement distillation (section 2.6.4) we do not require the conversion to be perfect. The only requirement is that the fidelity approaches unity as $n \rightarrow \infty$.

Note that to produce mixed states from Bell states using perfect LOCC we will need to discard some information. This can be done by doing a random local unitary transformation and not recording which one it was, or by doing a measurement and forgetting the outcome. A third option is to entangle the local system with an ancilla system and discard the ancilla. It is not surprising then that the distillable entanglement usually is strictly smaller than the entanglement cost. Actually, for any measure E which satisfies certain natural conditions for asymptotic measures, the entanglement cost and distillable entanglement provide upper and lower bounds [HHH00]²,

$$E_D \leq E \leq E_C \quad (4.11)$$

The bound entangled states, discussed in section 3.2.2, is an example where the difference between E_C and E_D is highly visible. E_D is obviously zero, but E_C is finite. This illustrates that one single measure is not enough to quantify the entanglement resources in a mixed quantum state, even in the asymptotic limit.

4.2.2 Axioms for abstract measures

The axiomatic approach to quantifying entanglement has been quite successful in finding measures that quantify certain aspects of entanglement. During the course of finding the right conditions, some conditions have been discarded as unnecessary and others have been restricted to a specific regime. The common denominator, though, has always been that entanglement, irrespective of how we quantify it, should not increase on average by any LOCC operation, i.e. it is monotonic under LOCC. The first attempt was done by Vedral *et al.* in [VPRK97] and was slightly improved by some of the same authors in [VP98]. Vidal [Vid00] argued that the only absolute requirement was monotonicity under LOCC. This automatically implies other properties such as nonchange under local unitary transformations, convexity and constancy on separable states.

The conditions on mixed states discussed here are taken from [DHR02]. The previously discussed conditions on pure states are specialisations of those.

(E0a) $E(\rho) = 0$ if and only if ρ is separable. This is a useful property for measures that satisfy it, but it is too strict in general. Bound entangled states, for instance, have zero distillable entanglement, but are not separable.

(E0b) $E(\rho) = 0$ if ρ is separable.

The normalisation conditions are the same as for pure states.

(E1a) (Normalisation) For the maximally entangled state in $d \times d$ dimensions, $|\Phi_d^+\rangle \langle \Phi_d^+|$,

²In [HHH00] the upper bound was E_F , the regularised *entanglement of formation*. This has been shown to be equal to the asymptotic entanglement cost [HHT01].

where

$$|\Phi_d^+\rangle = \sum_{i=1}^d \frac{1}{\sqrt{d}} |i\rangle_A \otimes |i\rangle_B \quad (4.12)$$

and the two sets $\{|i\rangle_A\}$ and $\{|i\rangle_B\}$ are orthonormal bases, $E(|\Phi_d^+\rangle \langle \Phi_d^+|) = \log d$.

(E1b) For a Bell state $|\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, $E(|\Phi^+\rangle \langle \Phi^+|) = 1$.

(E2a) (LOCC monotonicity) For any LOCC operation Λ , $E(\Lambda(\rho)) \leq E(\rho)$.

(E2b) When Λ is a strictly local operation which is either unitary or adds extra dimensions, then $E(\Lambda(\rho)) = E(\rho)$.

(E2c) When Λ is a strictly local unitary operation, then $E(\Lambda(\rho)) = E(\rho)$.

(E3a) (Continuity) Let $\{\rho_n\}$ and $\{\sigma_n\}$ be sequences of bipartite states living on the sequence of Hilbert spaces $\{\mathcal{H}_n\}$. For all such sequences such that $\|\rho_n - \sigma_n\|_1 \rightarrow 0$ where $\|\cdot\|_1$ is the trace norm $\|A\|_1 = \text{tr}(\sqrt{A^\dagger A})$,

$$\frac{E(\rho_n) - E(\sigma_n)}{1 + \log(\dim \mathcal{H}_n)} \rightarrow 0. \quad (4.13)$$

(E3b) The previous condition (E3a) is weakened by only requiring it to be satisfied for approximations to pure states. That is, (E3a) only needs to hold when the ρ_n are pure.

Many measures satisfy the weak additivity condition, but it is not considered necessary for a good measure.

(E4a) (Weak additivity) For all states ρ and $n \geq 1$,

$$\frac{E(\rho^{\otimes n})}{n} = E(\rho). \quad (4.14)$$

(E4b) (Asymptotic weak additivity) Given $\epsilon > 0$ and a state ρ , there exists an integer $N > 0$ such that for all $n \geq N$,

$$\frac{E(\rho^{\otimes n})}{n} - \epsilon \leq E(\rho) \leq \frac{E(\rho^{\otimes n})}{n} + \epsilon. \quad (4.15)$$

(E5a) (Subadditivity) For all states ρ and σ ,

$$E(\rho \otimes \sigma) \leq E(\rho) + E(\sigma). \quad (4.16)$$

(E5b) A special case of the above is when ρ and σ are different numbers of the same state tensored together. For all states ρ , and $m, n \geq 1$,

$$E(\rho^{\otimes(m+n)}) \leq E(\rho^{\otimes m}) + E(\rho^{\otimes n}). \quad (4.17)$$

(E5c) (Existence of a regularisation) For all bipartite states ρ , the limit

$$E^\infty(\rho) \equiv \lim_{n \rightarrow \infty} \frac{E(\rho^{\otimes n})}{n} \quad (4.18)$$

exists. E^∞ is called the regularisation of E . This is the weakest of the additivity conditions, and is usually satisfied.

(E6a) (Convexity) Mixing of states does not increase entanglement. For all bipartite states ρ and σ ,

$$E(\lambda\rho + (1 - \lambda)\sigma) \leq \lambda E(\rho) + (1 - \lambda)E(\sigma) \quad (4.19)$$

for $0 \leq \lambda \leq 1$.

Although (E6a) seems very natural as forgetting which of a set of prepared states you have should not increase entanglement, there are indications that the (asymptotic) distillable entanglement is not convex [SST01]. This is based on the fact that PPT bound entangled states together with NPPT bound entangled states (if they exist) can be distilled. Thus, even a mixture of two states with zero distillable entanglement can be distilled. A weaker condition which is shown in [DHR02] to hold also for the distillable entanglement is that the convexity need only hold on decompositions into pure states.

(E6b) For any bipartite state and any pure state realisation $\rho = \sum_i p_i |\psi_i\rangle \langle \psi_i|$, $p_i \geq 0$ and $\sum_i p_i = 1$,

$$E(\rho) \leq \sum_i p_i E(|\psi_i\rangle \langle \psi_i|). \quad (4.20)$$

Like for pure states, asymptotic weak additivity is sufficient for weak additivity, so (E4a) and (E4b) are actually equivalent. The same goes for the subadditivity conditions (E5a) and (E5b).

4.2.3 E_D and E_C as extreme measures

In [DHR02] the exact conditions for E_D and E_C to be the lower and upper bound for an entanglement measure were investigated. Three versions of the theorem were found. With different conditions on the measures either the nonregularised or the regularised measure would be bounded by E_D and E_C .

(i) For an entanglement measure E satisfying (E1a), (E2a), (E3a) and (E4b) (and thereby (E4a)), for all states ρ ,

$$E_D(\rho) \leq E(\rho) \leq E_C(\rho). \quad (4.21)$$

These conditions are very strong, and until recently no function was known that satisfied them. The “squashed entanglement” [CW04], however, satisfies all conditions.

(ii) For an entanglement measure E satisfying (E1a), (E2a), (E3a) and (E5c), then for all states ρ , the *regularised* version E^∞ (which exists by (E5c) and always satisfies (E4a), but not necessarily (E3a)) is bounded by E_D and E_C .

$$E_D(\rho) \leq E^\infty(\rho) \leq E_C(\rho) \quad (4.22)$$

These conditions are easier to satisfy, and they are satisfied by both the entanglement of formation and the relative entropy of entanglement.

(iii) Let E be an entanglement measure satisfying (E1a), (E2a), (E3b) and (E6b). Then if weak additivity (E4a) holds,

$$E_D(\rho) \leq E(\rho) \leq E_C(\rho) \quad (4.23)$$

and if subadditivity (E5a) holds,

$$E_D(\rho) \leq E^\infty(\rho) \leq E_C(\rho). \quad (4.24)$$

4.2.4 Relative entropy of entanglement and other distance based measures

Along with the first attempts to give realistic conditions for entanglement measures [VPRK97, VP98], a class of measures that satisfied the conditions was introduced. The conditions considered was that the measure would be zero if and only if the state is separable (E0a), that local unitary operations leave it constant (E2c) and that its expectation value does not increase under LOCC (E2a).

The class of measures is based on some distance function (but not necessarily a metric) on the set of density matrices. The measure is then the distance from the state in question to the nearest separable state. That is, for a distance function $D(\rho, \sigma)$ an entanglement measure corresponding to the distance function could be defined as

$$E(\rho) \equiv \inf_{\sigma \in \mathcal{S}(\mathcal{H})} D(\rho, \sigma) \quad (4.25)$$

where $\mathcal{S}(\mathcal{H})$ is the set of separable states on $\mathcal{H}$. The conditions imposed on E is now converted into conditions on D .

The quantum *relative entropy*³ can be used as a distance function on the set of density operators. It is defined as

$$S(\rho \parallel \sigma) \equiv \text{tr}[\rho(\log \rho - \log \sigma)]. \quad (4.26)$$

The relative entropy is not a metric and is not even symmetric. It is nonnegative, and zero only for identical density operators. The same unitary operation on both states leaves it invariant. The quantum relative entropy can be interpreted as a distinguishability of quantum states. More precisely, suppose we are given a large (but finite) number n of quantum systems that are all in the same state, which is either ρ or σ . Our task is to perform measurements to infer which. The probability for inferring from optimal measurements on the composite system, that the given state is ρ when it really is σ is [VPJK97]

$$P_n(\sigma \rightarrow \rho) = 2^{-nS(\rho \parallel \sigma)} \quad (4.27)$$

(for large n). This expression is not symmetric in ρ and σ .

To illustrate this, consider the qubit states

$$\rho = |0\rangle\langle 0| \quad \text{and} \quad \sigma = \frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1| = \frac{1}{2}\mathbb{I}_2$$

i.e. one arbitrary pure state (denoted $|0\rangle$) and the completely mixed state. These states are optimally distinguished by measurement if we measure each particle in the $\{|0\rangle, |1\rangle\}$ basis. In the case that we are performing measurements on σ , the probability for measuring $|0\rangle$ and $|1\rangle$ are both $\frac{1}{2}$. If all measurements give $|0\rangle$, we would wrongly conclude that the state was ρ . The probability for that to happen is 2^{-n} . Conversely, if we perform the measurements on systems in state ρ , we will always get the result $|0\rangle$, so the probability for confusing it with σ is zero.

³For reviews of the role of the relative entropy in quantum information theory, see [Ved02, SW02].

The quantum relative entropy can be calculated from the eigenvalues and eigenvectors of the density operators as shown in Appendix A. In the above example, it can be calculated from (A.4) giving $S(\rho\|\sigma) = 1$ and $S(\sigma\|\rho) = +\infty$. This is consistent with the probabilities above and (4.27). It is a general feature that $S(\sigma\|\rho) = +\infty$ when ρ is a pure state. This is because for any pure state there is a complete measurement for which the outcome is 100% certain, and using this measurement would make it impossible to confuse it with another state.

This property alone makes the relative entropy with the arguments reversed, $D(\rho, \sigma) = S(\sigma\|\rho)$, unusable in (4.25) as a distance function. It would give infinity for any pure entangled ρ . $S(\rho\|\sigma)$ on the other hand, behaves well for separable σ and the entanglement measure generated by it,

$$E_r(\rho) \equiv \inf_{\sigma \in \mathcal{S}(\mathcal{H})} S(\rho\|\sigma), \quad (4.28)$$

is called the *relative entropy of entanglement* [VP98]. It could be said that it quantifies the unlikelihood for a separable state to give measurement outcomes consistent with the entangled state.

From its definition and the fact that $S(\rho\|\sigma) = 0$ only when $\rho = \sigma$, it is obvious that the relative entropy of entanglement satisfies (E0a) ($E_r(\rho) = 0$ iff $\rho \in \mathcal{S}(\mathcal{H})$). It reduces to the entropy of entanglement on pure states [VP98], and therefore satisfies the normalisation criterion (E1a) and continuity on pure states (E3b) by the uniqueness theorem. It is nonincreasing under LOCC operations (E2a) [VPRK97]. Continuity (E3a) was shown in [DH99]. There have been found counterexamples to weak additivity (E4a) [VW01] and thereby also to asymptotic weak additivity (E4b). Actually, for very high dimension of a certain class of states, $E_r(\rho) \approx E_r(\rho \otimes \rho)$. It can also be shown that it is both subadditive (E5a) and convex (E6a) [DHR02].

The regularisation (E5c)

$$E_R(\rho) \equiv E_r^\infty = \lim_{n \rightarrow \infty} \frac{E_r(\rho^{\otimes n})}{n} \quad (4.29)$$

of the relative entropy of entanglement satisfies automatically (E0a), (E1a), (E2a), (E5a) and (E6a) as those conditions extend to any regularisation of a measure that satisfies them [DHR02]. In addition, as any regularisation does, it satisfies weak additivity (E4a).

Any distance based measure, including the relative entropy of entanglement, can be varied by considering other classes of states than the separable ones. What is important is that the class is closed under LOCC. For instance, the set can be extended to PPT states. This will give a measure which is easier to calculate and inferior or equal to the version for separable states. For 2×2 and 2×3 dimensional systems the measures coincide as the PPT is then equivalent to separability.

Even though the quantum relative entropy is by far the most used of the distance based measures, other distance functions provide usable entanglement measures as well. Another measure of distinguishability of quantum states is the *Bures metric* [Bur69, Hüb92, FC95], in this context defined as [VP98]

$$D_B(\rho\|\sigma) \equiv 2 - 2\sqrt{F(\rho, \sigma)} \quad (4.30)$$

where

$$F(\rho, \sigma) \equiv \left[\text{tr} \left\{ (\sqrt{\sigma} \rho \sqrt{\sigma})^{1/2} \right\} \right]^2 \quad (4.31)$$

is called Uhlmann's transition probability [Uhl76]. The Bures metric is a true metric and thus symmetric. It is a generalisation of the pure state Fubini-Study metric to mixed states [Hüb92]. The entanglement measure generated by it satisfies (E0a), (E1b) and (E2a). Unlike the relative entropy of entanglement, however, it doesn't reduce to the entropy of entanglement on pure states. Actually, it is smaller than the entropy of entanglement [VP98].

There may be other distance functions that generate entanglement measures with nice properties, but the two mentioned are the ones used in practice. The Hilbert-Schmidt distance

$$D_{HS}(\rho, \sigma) \equiv \|\rho - \sigma\|_{HS}^2 = \text{tr} [(\rho - \sigma)^2] \quad (4.32)$$

was suggested as a candidate [VP98, WT99]. However, it was shown by Ozawa [Oza00] that the distance violated one of the criteria it was conjectured to have (nonexpansion under physical operations), so it did not satisfy the sufficient conditions for it to be nonincreasing under LOCC. On the other hand, the very similar trace norm distance

$$D_T(\rho, \sigma) \equiv \|\rho - \sigma\|_1 = \text{tr} \left[\sqrt{(\rho - \sigma)^2} \right] \quad (4.33)$$

has been shown to generate an entanglement measure which is monotonic under LOCC [EAP03].

4.2.5 Entanglement of formation

The *entanglement of formation* was historically the first entanglement measure to appear [BDSW96]. In the literature it is occasionally called *entanglement of creation*. It was meant to be the asymptotic entanglement cost, but this interpretation rests on the weak additivity which is strongly conjectured, but still not proved. However, the regularisation of it has been shown to be equal to the asymptotic entanglement cost [HHT01].

The entanglement of formation is a straightforward generalisation of the entropy of entanglement to mixed states. Remember that in the asymptotic limit, the entanglement cost in Bell states of preparing a pure state is given by the entropy of entanglement. Thus, it was natural to define the entanglement of formation for a pure state as the entropy of entanglement. For a given ensemble of pure states $\mathcal{E} = \{p_i, |\psi_i\rangle\}$ the entanglement of formation is the average of the entropy of entanglement for the states in the ensemble

$$E_f(\mathcal{E}) \equiv \sum_i p_i E_E(|\psi_i\rangle). \quad (4.34)$$

A mixed state can be realised by a multitude of pure state ensembles, with different entanglement of formation. As any of those ensembles realises the mixed state, the natural definition for the entanglement of formation for a mixed state is the entanglement of formation for the “most economic” ensemble. That is, the entanglement of formation for a mixed state is defined as

$$E_f(\rho) \equiv \inf_{\mathcal{E}} \sum_i p_i E_E(|\psi_i\rangle) \quad (4.35)$$

where the infimum is taken over all ensembles $\varepsilon = \{p_i, |\psi_i\rangle\}$ that realises the state ρ .

How come entanglement of formation is not necessarily equal to the entanglement cost? It is easy to see that the asymptotic entanglement cost cannot be greater than the entanglement of formation. Imagine that we are to produce a large number n_{out} ($n_{\text{out}} \rightarrow \infty$) of some bipartite state ρ_{out} . We find the pure state realisation $\{p_i, |\psi_i\rangle\}$ of ρ_{out} that has the lowest average entropy of entanglement (equal to the entanglement of formation). Then we produce this ensemble by producing the pure states in amounts corresponding to the probability distribution $\{p_i\}$. In the asymptotic limit this can be done reversibly. When we mix the produced states (discard the information saying which is which), we have a collection of systems in the state ρ_{out} . Since the entropy of entanglement is conserved in the first step, and $E_E = 1$ for Bell states, we have produced systems in state ρ_{out} at a cost equal to the entanglement of formation.

The question is if we can do better than this. Intuitively the above is the most economical way, as any other realisation of ρ_{out} would give a higher or equal cost. But there is another way of producing a large number of ρ_{out} . We can produce n_{out}/N copies of the state $\rho_{\text{out}}^{\otimes N}$. It is not known whether the optimal pure state realisation of this state will have an average entropy of entanglement N times that of the state ρ_{out} . In other words it is not known if E_f is weakly additive, $E_f(\rho) = \frac{1}{N}E_f(\rho^{\otimes N})$. If it is, then $E_f = E_F \equiv E_f^\infty$ which is equal to the asymptotic entanglement cost E_C . Additivity has been shown for some special classes of states [BN01, VW01, VDC02, Shi02], but whether it holds in general, is one of the big open questions in quantum information theory.

From the definition of E_f , it is easy to see that it satisfies (E0a) and (E1a). (E0a) by the fact that the optimal pure state decomposition of a separable states is into a mixture of product states, (E1a) by the fact that it reduces to the entropy of entanglement on pure states. LOCC monotonicity (E2a) was shown when it was introduced [BDSW96], and continuity (E3a) was shown in [Nie00]. Subadditivity (E5a) follows from the discussion of additivity above, one may use the same type of decomposition for a tensor product, but one *might* be able to do better, which would give a lower E_f . Convexity (E6a) follows more or less directly from the definition.

The way the entropy of entanglement was extended to mixed states in the case of the entanglement of formation, can also be used for other measures on pure states. Measures constructed this way are called *convex roof measures* [Uhl98, Hor01]. The method extends a measure of some set (here pure states) to the convex hull⁴ (here mixed states), where it is the largest function that is convex and compatible with the measure on the original set. The method was used in [LCOK03] construct an entanglement measure that extends the entanglement measure called *negativity* to mixed states in another way than the original measure.

A concept related to the entanglement of formation is the *concurrence* [HW97, Woo98]. It is defined for a system of two qubits. For a general state ρ of two qubits, let $\tilde{\rho}$ be the spin-flipped state

$$\tilde{\rho} \equiv (Y \otimes Y)\rho^*(Y \otimes Y) \quad (4.36)$$

where the Y is the Pauli Y operator and ρ^* is the complex conjugate of ρ , both taken in

⁴The convex hull of a set is the set of all elements that can be written as a convex combination of the original set. For instance, the convex hull of two points is the line connecting them.

the standard basis $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$. Let the Hermitian matrix R be defined as

$$R \equiv \sqrt{\sqrt{\rho}\tilde{\rho}\sqrt{\rho}} \quad (4.37)$$

with eigenvalues in decreasing order $\{\lambda_i\}$. The concurrence is then defined as

$$C(\rho) \equiv \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\}. \quad (4.38)$$

The concurrence is monotonic under LOCC, and can thus be used as an entanglement measure for two qubits. The great advantage is that it is easily computable. But more important is that it is directly related to the entanglement of formation, providing an explicit formula for the entanglement of formation in the case of two qubits. Let the function $\mathcal{E}$ be given by

$$\mathcal{E}(C) \equiv h\left(\frac{1 + \sqrt{1 - C^2}}{2}\right) \quad (4.39)$$

where h is the binary entropy function

$$h(x) \equiv -x \log x - (1 - x) \log(1 - x). \quad (4.40)$$

Then the entanglement of formation is simply given by [HW97, Woo98]

$$E_f(\rho) = \mathcal{E}(C(\rho)). \quad (4.41)$$

There have been various attempts to generalise the concept of concurrence. Uhlmann [Uhl00] considered general conjugations instead of the special (4.36). This was further generalised to concurrence vectors by Audenaert *et al.* [AVD01]. The previously mentioned convex roof extended negativity coincides with the concurrence in the case of two qubits [LCOK03], and thus provides another generalisation. Some of those generalisations – along with other aspects of concurrence and entanglement of formation – are reviewed in [Woo01].

4.2.6 Negativity

The entanglement measures discussed so far have all had a serious drawback. Their definition includes some kind of optimisation, which make their evaluation very difficult. In fact, expressions for the relative entropy of entanglement and entanglement of formation has only been calculated for highly symmetric states (e.g. [VP98, TV00, VW01, ADVW02]) and low dimensional cases (e.g. [HW97, Woo98]). For general states, evaluating a measure includes heavy numerical calculations. With the aim of introducing a *computable* measure of entanglement, two related quantities were defined by Vidal and Werner [VW02a]. They can both be seen as a quantification of the PPT criterion for separability, and because of that they have the disadvantage that they fail to distinguish between separable states and entangled PPT states. Both quantities are based on the trace norm of the partial transpose (in some basis) of a state, $\|\rho^{T_B}\|_1$, which can easily be calculated using standard linear algebra packages. The first quantity is the *negativity*

$$\mathcal{N}(\rho) \equiv \frac{\|\rho^{T_B}\|_1 - 1}{2}. \quad (4.42)$$

This quantity is equal to the absolute value of the sum of negative eigenvalues of ρ^{T_B} . It was first introduced in [ŽHSL98, Appendix B] (without the factor $\frac{1}{2}$) and later shown to not increase under LOCC [VW02a]. The other quantity is the *logarithmic negativity*

$$E_{\mathcal{N}}(\rho) \equiv \log \|\rho^{T_B}\|_1, \quad (4.43)$$

which is not strictly monotonic under LOCC, but does not increase for a subclass of LOCC.

For all separable states (and more generally all PPT states), both $\mathcal{N}$ and $E_{\mathcal{N}}$ vanish. This is easily seen from the fact that the partial transpose operation does not change the trace of the density operator. If ρ^{T_B} is a positive operator (i.e. ρ satisfies the PPT criterion for separability), then $\|\rho^{T_B}\|_1 = 1$, since $\|\rho^{T_B}\|_1$ is the sum of the absolute values of the eigenvalues of ρ^{T_B} and all eigenvalues are positive. For states violating the PPT criterion, $\|\rho^{T_B}\|_1 > 1$. In this sense, both quantities quantify how much the state violates the PPT criterion.

A lot of properties of $\mathcal{N}$ and $E_{\mathcal{N}}$ were deduced in [VW02a]. As already mentioned, $\mathcal{N}$ is zero for all separable states, but is also zero for PPT entangled states. Hence, it satisfies (E0b) but not (E0a). It is $\frac{1}{2}$ for Bell states, so it doesn't satisfy the normalisation criteria either. It could be made to satisfy (E1b) by defining it as the double value, but the normalisation wouldn't extend to maximally entangled states in dimension $d \times d$ (E1a). It is monotonic under LOCC (E2a), convex (E6a), and superadditive, satisfying $\mathcal{N}(\rho \otimes \sigma) = \mathcal{N}(\rho) + \mathcal{N}(\sigma) + 2\mathcal{N}(\rho)\mathcal{N}(\sigma)$. Thus, neither additivity of any kind (E4a) nor subadditivity (E5a) is satisfied.

Like $\mathcal{N}$, logarithmic negativity $E_{\mathcal{N}}$ satisfies (E0b) and not (E0a). However, unlike $\mathcal{N}$ it also satisfies the normalisation condition (E1a). But it does not coincide with the entropy of entanglement for pure states in general. Actually $E_{\mathcal{N}} \geq E_E$ on pure states, with equality only for maximally entangled states. It is not monotonic under LOCC (E2a), but is an upper bound for the distillable entanglement. Nor is it convex (E6a). The most appealing property of $E_{\mathcal{N}}$ is probably that it satisfies (strong) additivity, $E_{\mathcal{N}}(\rho \otimes \sigma) = E_{\mathcal{N}}(\rho) + E_{\mathcal{N}}(\sigma)$. From this, weak additivity (E4a) and subadditivity (E5a) comes for free.

The negativity can be generalised to give rise to other entanglement measures. This is done by replacing $\|\rho^{T_B}\|_1$ by another norm $\|\rho\|$. The generalisation comes naturally if we define the trace norm in another, but equivalent way. Any Hermitian matrix can be written as a difference of two positive operators,

$$A = a_+ \rho^+ - a_- \rho^- \quad (4.44)$$

where $\rho^{\pm}$ are density matrices and $a_{\pm}$ are nonnegative numbers. There exists a decomposition of the form (4.44) which minimises $a_+ + a_-$. For this decomposition the trace norm is $\|A\|_1 = a_+ + a_-$, and a_- is the absolute sum of the negative eigenvalues of A [VW02a]. From this we get another expression for the negativity, namely

$$\mathcal{N}(A) = \inf\{a_- \mid A^{T_B} = a_+ \rho^+ - a_- \rho^-\}. \quad (4.45)$$

Minimising a_- is in this case the same as minimising $a_+ + a_-$, since by taking the trace of the condition, we get $a_+ = a_- + \text{tr}(A^{T_B})$. If A^{T_B} is a density matrix, A is PPT, and we can take a_- to be zero, so the negativity vanishes as it should for PPT states.

The condition $A^{TB} = a_+\rho^+ - a_-\rho^-$ restricts the set of density matrices $\rho^\pm$ over which we can take the infimum. Another norm and another negativity can be obtained by taking another set S from which $\rho^\pm$ must be picked. Norms that can be defined in this way corresponding to a compact set S are called *base norms*. The norm corresponding to a general compact set S is then

$$\|A\|_S = \inf\{a_+ + a_- \mid A = a_+\rho^+ - a_-\rho^-, a_\pm \geq 0, \rho^\pm \in S\} \quad (4.46)$$

and the corresponding negativity

$$\mathcal{N}_S(A) = \inf\{a_- \mid A = a_+\rho^+ - a_-\rho^-, a_\pm \geq 0, \rho^\pm \in S\}. \quad (4.47)$$

We get back the usual trace norm and negativity from this by taking S as the set of all Hermitian (not necessarily positive) matrices with unit trace such that the partial transpose is positive. In mathematical terms, $S = \{A \mid A = A^\dagger, \text{tr}(A) = 1, A^{TB} \geq 0\}$.

By choosing S as the set of all separable states, another entanglement measure arises. It was described prior to the introduction of negativity and was introduced in [VT99]. It is called *robustness of entanglement* and was defined as the minimum amount of mixing with locally prepared states needed to make the state separable. Also other measures have been related to negativities [Hor01], such as the measures introduced by Rudolph based on the greatest cross norm [Rud01]. It should be noted that only the original negativity has the advantage of being easily computable. The other negativities need to be evaluated by finding the infimum over a large parameter space.

4.2.7 Squashed entanglement

Of the entanglement measures considered so far, only few are additive, and when they are some of the other greatly desired conditions are not fulfilled. Quite recently a new entanglement measure was proposed, having more nice properties than its predecessors. It is motivated from the so called intrinsic information of classical cryptography and is called *squashed entanglement* [CW04]. It is defined as

$$E_{\text{sq}}(\rho) \equiv \inf \left\{ \frac{1}{2} I(A; B|E) \mid \rho^{ABE} \text{ extension of } \rho \text{ to } \mathcal{H}_E \right\} \quad (4.48)$$

The infimum is taken over all extensions to a third subsystem E , so $\rho \equiv \rho^{AB} = \text{tr}_E(\rho^{ABE})$. $I(A; B|E)$ is the quantum mutual conditional information defined as

$$I(A; B|E) \equiv S(\rho^{AE}) + S(\rho^{BE}) - S(\rho^{ABE}) - S(\rho^E). \quad (4.49)$$

Most of the properties of E_{sq} were shown along with its introduction as an entanglement measure in [CW04]. The squashed entanglement vanishes on all separable states (E0b), but it is not known whether it also vanishes for some entangled states (E0a). If there exists an extension to $\mathcal{H}_E$ with $\dim \mathcal{H}_E < \infty$ for which $I(A; B|E)$ vanishes, it implies that ρ^{AB} is separable. But the infimum in (4.48) is taken over all extensions, and thus $E_{\text{sq}}(\rho^{AB})$ may be zero even if no finite dimensional extension makes $I(A; B|E)$ zero. It coincides with the entropy of entanglement on pure states, and thus satisfies the normalisation condition

(E1a). It is also bounded from below by the distillable entanglement and from above by the entanglement cost. As a proper entanglement measure it does not increase under LOCC (E2a) and it is convex (E6a). When it was introduced, it was shown to be continuous on most of the state space, and indications that it was continuous on the whole state space were presented. Continuity was finally proved by Alicki and Fannes [AF04] (E3a). The measure is strongly additive, making it satisfy both weak additivity (E4a) and subadditivity (E5a). So except from (E0a), which is still unknown, squashed entanglement satisfies the strictest versions of all conditions presented.

In [VW01] various additivity properties were discussed. The additivities we have presented so far are for *independent* pairs, described by tensor products. But if there are entanglement or classical correlations between the pairs as well, they may be taken advantage of. Consider the state $\rho^{ABA'B'}$ which lives on the Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$, where $\mathcal{H}_A \otimes \mathcal{H}_{A'}$ is controlled by Alice and $\mathcal{H}_B \otimes \mathcal{H}_{B'}$ by Bob. Each of the primed and unprimed subsystems are individually described by tracing out the other,

$$\rho^{AB} = \text{tr}_{A'B'}(\rho^{ABA'B'}), \quad \rho^{A'B'} = \text{tr}_{AB}(\rho^{ABA'B'}), \quad (4.50)$$

and if they are independent $\rho^{ABA'B'} = \rho^{AB} \otimes \rho^{A'B'}$. For a strongly additive entanglement measure this implies $E(\rho^{ABA'B'}) = E(\rho^{AB}) + E(\rho^{A'B'})$. But if there are correlations between the pairs that can be exploited, we would expect

$$E(\rho^{ABA'B'}) \geq E(\rho^{AB}) + E(\rho^{A'B'}). \quad (4.51)$$

This property is called *strong superadditivity* [VW01] and is satisfied by the squashed entanglement [CW04].

Like most other entanglement measures, the definition of squashed entanglement involves an optimisation which makes the measure hard to compute. Due to its recent introduction, not much work has been published about its calculation on special sets of states, nor about efficient numerical algorithms. One of the important open questions is whether it vanishes on any entangled states, especially on the bound entangled ones.

4.2.8 Other entanglement measures

In addition to the entanglement measures discussed, other measures have been proposed without gaining the same importance. One of them is the *entanglement of assistance* [Coh98, DFM⁺99]. It is based on the following idea. Many copies of an entangled tripartite pure state, $|\Psi^{ABC}\rangle$, is shared between Alice, Bob and Charlie. Now, Alice and Bob want to use the state to perform some task involving the use of bipartite entanglement. If they ignore Charlie completely, they share the state $\rho^{AB} = \text{tr}_C(|\Psi^{ABC}\rangle \langle \Psi^{ABC}|)$. In other words they share a mixed state, and it might even be separable. But Charlie wants to help, and Alice and Bob may communicate classically with him. Therefore, Charlie can perform local operations on his part of the system and tell Alice and Bob about the measurement outcomes. A measurement by Charlie will then leave the subsystem of Alice and Bob in some pure state, which depending on the measurement outcome may or may not be entangled. Alice and Bob can then discard the pairs that are not entangled, and turn the rest reversibly (in the asymptotic limit) into Bell states. If each measurement by Charlie leaves them with a

state $|\Psi_i^{AB}\rangle$ with probability p_i , the number of Bell states produced per copy of the tripartite state will be equal to the average entropy of entanglement of the ensemble $\mathcal{E} = \{p_i, |\Psi_i^{AB}\rangle\}$, namely

$$E(\mathcal{E}) = \sum_i p_i E_E(|\Psi_i^{AB}\rangle). \quad (4.52)$$

Now, Charlie wants to do the measurements in a way that leaves Alice and Bob with as much entanglement as possible on average. It turns out that the optimisation can be taken over all ensembles $\mathcal{E} = \{p_i, |\Psi_i^{AB}\rangle\}$ consistent with ρ^{AB} . Thus, what Charlie can achieve does not depend on the initial tripartite state $|\Psi^{ABC}\rangle$, only on ρ^{AB} , so all pure tripartite states for which ρ^{AB} is the same, allow Charlie to help Alice and Bob to the same result. The entanglement of assistance is the average entropy of entanglement Alice and Bob are left with after Charlie has applied his optimal strategy, namely

$$E_a(\rho^{AB}) \equiv \sup_{\mathcal{E}} \sum_i p_i E_E(|\Psi_i^{AB}\rangle). \quad (4.53)$$

We can see that this is dual to the entanglement of formation (4.35), which is the same with an infimum instead of the supremum. Even though the entanglement of formation is believed to be additive, there are states for which the entanglement of assistance is superadditive,

$$E_a(\rho^{AB} \otimes \sigma^{AB}) \geq E_a(\rho^{AB}) + E_a(\sigma^{AB}). \quad (4.54)$$

A recently introduced measure related to the entanglement of assistance is the *localizable entanglement* [VPC04]. It is really a multipartite measure for a large number of qubits, but the goal is, just as for the entanglement of assistance, to optimise (localise) the entanglement between two subsystems by performing measurements on the others. If we had allowed global measurements on the other qubits, we could have regarded it as a single ‘‘Charlie’’ system, and it would have reduced to the entanglement of assistance. Because we can only perform local measurements on each individual qubit, the localizable entanglement is necessarily smaller than the entanglement of assistance.

Another entanglement measure recently introduced is the *witnessed entropy of entanglement* [BV04]. It is based on the concept of entanglement witnesses [HHH96]. An entanglement witness for an entangled state σ is a Hermitian operator W for which $\text{tr}(W\sigma) < 0$ and $\text{tr}(W\rho) \geq 0$ for all separable states ρ . We conventionally normalise entanglement witnesses to have $\text{tr}(W) = 1$. We say that an entanglement witness W_σ is optimal for a state σ if

$$\text{tr}(W_\sigma\sigma) \leq \text{tr}(W\sigma) \quad (4.55)$$

for all entanglement witnesses W . The definition of the witnessed entropy of entanglement is based on the optimal witness,

$$E_w(\rho) \equiv \log(D/d) \max[0, -\text{tr}(W_\rho\rho)] \quad (4.56)$$

where D is the dimension of the total Hilbert space and d the smallest of the dimensions of the Hilbert spaces of the subsystems. Note that the definition applies equally well to bipartite and multipartite systems. It is not additive, but satisfies the rest of our conditions for entanglement measures, (E0a), (E1a), (E2a), (E3a), (E5a) and (E6a). The main

advantage of this measure is that it can be approximately calculated for all mixed states. Finding the exact optimal entanglement witness, is a hard optimisation problem, but using approximation techniques, there are algorithms to find the E_w to a given precision.

Other measures can be defined operationally in the following way. Take your favourite application of entanglement, study how well or bad it performs for different sets of states, define an entanglement measure from this and check that it satisfies at least some of the conditions for entanglement, above all LOCC monotonicity. One of the operational measures that have had limited impact on the community is an entanglement measure introduced by Hiroshima [Hir02] based on the capacity of dense coding [BW92]. Another operational measure by Biham *et al.* [BNO02] is derived from Grover's search algorithm [Gro96, Gro97]. It is based on the fact that the algorithm performs worse the more the input state is entangled.

Chapter 5

Beyond bipartite entanglement and finite dimension

The systems we have considered until now have all been bipartite systems with finite dimensional Hilbert spaces. Some of the concepts and measures considered are valid also in the multipartite systems and for entanglement of continuous variables. Others lose their meaning in this more general setting. In this chapter we consider some of the steps that need to be taken to generalise the concepts from the previous sections to multipartite systems and systems with continuous variables.

5.1 Multipartite systems

While the basic structure of bipartite entanglement is well understood, multipartite entanglement is still an active field of research. In bipartite systems the pure state entanglement in the asymptotic regime is well understood, since the von Neumann entropy of the reduced density matrices can be conserved under transformations. In general the Schmidt decomposition cannot be extended to multipartite systems [Per95], so the eigenvalues of the reduced density matrices of each system need not be the same, nor do the von Neumann entropies.

The questions about entanglement literally acquire new dimensions when more than two subsystems are considered. Between two parties all relations are between those two parties, and in terms of quantum communication there is only one single channel. Once we increase the number of parties to three, we have three pair relations in the system. Still, those three pair relations do not describe all correlations in the system (neither classical nor quantum). Consider for instance the pure tripartite entangled state called the GHZ¹ state

$$|GHZ\rangle \equiv \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle). \quad (5.1)$$

If we trace out any one of the systems, we get the density matrix shared by the two remaining parties

$$\rho = \frac{1}{2}(|00\rangle\langle 00| + |11\rangle\langle 11|) \quad (5.2)$$

¹Daniel M. Greenberger, Michael A. Horne and Anton Zeilinger

which is separable. Thus, the state contains no bipartite entanglement between the systems, but the tripartite entanglement is maximal.

5.1.1 Separability

Some notions can be taken directly over from bipartite systems, though. The generalisation of separability from (3.3) is straightforward. A state is separable if and only if it can be written as a convex combination of product states;

$$\sum_i p_i \rho_i^{(A)} \otimes \rho_i^{(B)} \otimes \cdots \otimes \rho_i^{(N)} \quad (5.3)$$

with $p_i \geq 0$ and $\sum_i p_i = 1$.

The classification of states does not end with this. An entangled multipartite state may or may not still be entangled after one of the subsystems are ignored (i.e. traced out). A multipartite state that loses its entanglement when any of the subsystems is traced out, is called *multiseparable* [Tha99].

The PPT criterion for separability trivially extends to the multipartite case. If a multipartite state is separable, then transposing any number of the subsystems must also give a valid (i.e. positive) density operator.

5.1.2 Measures of entanglement

The definition of entanglement cost and the distillable entanglement is based on a standard bipartite state, namely the Bell state. This is a bipartite state, so it is not obvious how to extend these measures to the multipartite case. Nielsen [Nie01] argued that this could be overcome by generalising the definition from Bell states to a definition in terms of qubits transferred (in the case of creation) or the number of CNOT gates that could be implemented (in the case of distillation). For instance, to create a Bell state it is necessary to transfer one qubit. Likewise the transfer of one qubit can be simulated by teleporting a state, spending one Bell state. If the entanglement cost is defined in terms of how many qubits that need to be transferred, it will coincide with the usual definition for bipartite states.

Most of the conditions we imposed on measures in section 4.2.2 generalise to the multipartite case. The only possible exceptions are of course the normalisation conditions (E1a) and (E1b) which are based on Bell states. Those would be unnatural to impose on multipartite entanglement, but it should be possible for a measure to satisfy them for states where only two of the subsystems are entangled.

The relative entropy of entanglement and other distance based measures do not have any reference to the bipartite case in their definition. Therefore, they are just as valid in the multipartite case as in the bipartite case.

The entanglement of formation suffers the same problem as the entanglement cost, and can be generalised in the same way. For a pure state we define the entanglement of formation as the number of qubits we need to transfer in order to prepare it (LOCC comes for free). Then we extend the definition to the mixed states by the convex roof method as usual. Another more formal generalisation was suggested by Wang [Wan01].

The meaning of entanglement measures in the multipartite setting is more vague than in the bipartite. In the bipartite setting it quantifies in some way how much that can be achieved by the two parties together. In the multipartite case there are more possibilities. For instance there may be some entanglement between all of the parties, as in the state

$$|W\rangle \equiv \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle) \quad (5.4)$$

which on tracing out one of the subsystems gives the density matrix

$$\rho = \frac{1}{3} |00\rangle \langle 00| + \frac{2}{3} |\Psi^+\rangle \langle \Psi^+| \quad (5.5)$$

which is an entangled state with negativity $\mathcal{N} = 0.206$ and entanglement of formation $E_f = 0.550$ (calculated from the expressions in sections 4.2.6 and 4.2.5). Or there may be no entanglement between the pairs, as in the GHZ state mentioned earlier. Both states can be prepared by sending two qubits (after one of the parties has created the state locally). While the different bipartite measures place different orderings on the states [VP00], ordering is even more ambiguous for multipartite states. The types of entanglement are qualitatively more different, and a state being very much entangled in one sense (as defined by one measure), can have very little or no entanglement in another sense. All this says is that, since a multipartite system is a more complicated structure, we cannot expect one single parameter to be as good a description of the entanglement in it as it is in the bipartite case.

5.2 Continuous variables

Most of quantum information theory has been formulated in the context of finite dimensional Hilbert spaces. However, from an experimental point of view, entanglement is just as interesting in the continuous variable regime. This has led to efforts to quantify entanglement also in the case where the Hilbert spaces have infinite dimension. Two papers by Eisert *et al.* [ESP02, EP03] summarise many of the new problems that arise and their possible solutions.

First of all, when the dimension of the Hilbert space is infinite, we have states with an infinite amount of entanglement. This should come as no surprise, as the entropy of entanglement for a maximally entangled state in $d \times d$ dimensions is $\log d$ (c.f. criteria (P1a) and (E1a)). What is more worrying is that the set of pure states with infinite entropy of entanglement is actually dense in the trace norm on the set of pure states [ESP02]. This means that arbitrarily close to any product state is a state which has infinite entanglement. The solution to this problem is to consider only states where the mean energy is bounded from above. This is a reasonable physical assumption, and on this subset of states the entropy of entanglement is continuous. Other measures can also be defined on this subset without exhibiting strange behaviour.

The definition of separability in a continuous variable bipartite system must be slightly altered from the one in section 3.2.1. The expression is still the same as (3.3), namely

$$\sum_i p_i \rho_i^{(A)} \otimes \rho_i^{(B)} \quad (5.6)$$

with p_i is a probability distribution. But in infinite dimension we can only require that the state can be approximated arbitrarily well by an expression of the form (5.6).

The PPT criterion for separability is also valid in infinite dimension and for the special class of so-called Gaussian states, it is also sufficient for separability [Sim00].

It is a well known fact in the finite dimensional case that there is a neighbourhood around the completely mixed state $\mathbb{I}/d$ where all states are separable [ŽHSL98, PR00, Myr04]. However, the volume of the separable ball diminishes as the dimension of the system grows. In the infinite dimensional setting, even when the energy is bounded from above, it has been shown [ESP02] that arbitrarily close to any state (in the trace norm), there is an entangled state.

The class of so-called Gaussian states has a special role in entanglement theory of continuous variables. These are states with a Gaussian Wigner function (a quasi-probability distribution in phase space which completely describes a quantum state). These states are often encountered in the laboratory, and they can be completely described by two parameters. For Gaussian states, and operators that takes Gaussian states to Gaussian states, many of the notions from finite dimensional entanglement theory have similar counterparts. See e.g. [EP03, KGLC03, GWK⁺03, WGK⁺04, RE04].

Appendix A

Expressions for the quantum relative entropy

The quantum relative entropy is defined as

$$S(\rho\|\sigma) \equiv \text{tr}\{\rho \log \rho - \rho \log \sigma\}. \quad (\text{A.1})$$

We want to calculate it in terms of the eigenvectors and eigenvalues of ρ and σ . We denote the eigenvalues of ρ (σ) by r_i (s_i) and its eigenvectors by $|r_i\rangle$ ($|s_i\rangle$). In the case of degenerate eigenvalues we assume that the eigenvectors have been properly orthogonalised.

The linearity of the trace gives

$$S(\rho\|\sigma) = \text{tr}\{\rho \log \rho\} - \text{tr}\{\rho \log \sigma\}$$

where the first term can be identified as $-S(\rho)$, where $S(\cdot)$ is the von Neumann entropy. This depends only on the eigenvalues of ρ :

$$\text{tr}\{\rho \log \rho\} = \sum_i r_i \log r_i.$$

The second term is more complicated as the eigenvectors of the density operators in general are not the same. We decompose the density operators using the spectral decomposition.

$$\begin{aligned} \rho \log \sigma &= \left(\sum_i r_i |r_i\rangle \langle r_i| \right) \log \left(\sum_j s_j |s_j\rangle \langle s_j| \right) \\ &= \left(\sum_i r_i |r_i\rangle \langle r_i| \right) \left(\sum_j \log(s_j) |s_j\rangle \langle s_j| \right) \\ &= \sum_{ij} r_i \log(s_j) \langle r_i | s_j \rangle |r_i\rangle \langle s_j| \end{aligned}$$

Then we decompose $|r_i\rangle$ in the eigendirections of σ , $|r_i\rangle = \sum_k \langle s_k | r_i \rangle |s_k\rangle$.

$$\rho \log \sigma = \sum_{ijk} r_i \log(s_j) \langle r_i | s_j \rangle \langle s_k | r_i \rangle |s_k\rangle \langle s_j|$$

Now we can take the trace of this to get the second term

$$\text{tr}\{\rho \log \sigma\} = \sum_{ij} r_i \log s_j |\langle r_i | s_j \rangle|^2$$

We can now put the two terms together to get the expression we wanted;

$$S(\rho||\sigma) = \sum_i r_i \left\{ \log r_i - \sum_j \log s_j |\langle r_i | s_j \rangle|^2 \right\}. \quad (\text{A.2})$$

This may also be written as the single sum

$$S(\rho||\sigma) = \sum_i \left\{ r_i \log r_i - \langle s_i | \rho | s_i \rangle \log s_i \right\}. \quad (\text{A.3})$$

In the case when the eigenvectors for the two density matrices are the same (i.e. ρ and σ commute), we can arrange the eigenvectors so that $\langle r_i | s_j \rangle = \delta_{ij}$. (A.2) then reduces to

$$S(\rho||\sigma) = \sum_i r_i (\log r_i - \log s_i). \quad (\text{A.4})$$

References

- [ADR82] A. Aspect, J. Dalibard and G. Roger, *Experimental Test of Bell's Inequalities Using Time-Varying Analyzers*, Phys. Rev. Lett. **49**(25), 1804–1807 (1982).
- [ADVW02] K. Audenaert, B. De Moor, K. G. H. Vollbrecht and R. F. Werner, *Asymptotic relative entropy of entanglement for orthogonally invariant states*, Phys. Rev. A **66**, 032310 (2002), quant-ph/0204143.
- [AF04] R. Alicki and M. Fannes, *Continuity of quantum conditional information*, J. Phys. A: Math. Gen. **37**, L55–L57 (2004), quant-ph/0312081.
- [AVD01] K. Audenaert, F. Verstraete and B. De Moor, *Variational characterizations of separability and entanglement of formation*, Phys. Rev. A **64**, 052304 (2001), quant-ph/0006128.
- [BBC⁺93] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres and W. K. Wootters, *Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels*, Phys. Rev. Lett. **70**, 1895–1899 (1993).
- [BBP⁺96] C. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. Smolin and W. Wootters, *Purification of Noisy Entanglement and Faithful Teleportation via Noisy Channels*, Phys. Rev. Lett. **76**(5), 722–725 (1996), quant-ph/9511027.
- [BBPS96] C. Bennett, H. Bernstein, S. Popescu and B. Schumacher, *Concentrating partial entanglement by local operations*, Phys. Rev. A **53**(4), 2046–2052 (1996), quant-ph/9511030.
- [BDSW96] C. Bennett, D. DiVincenzo, J. Smolin and W. Wootters, *Mixed-state entanglement and quantum error correction*, Phys. Rev. A **54**(5), 3824–3851 (1996), quant-ph/9604024.
- [Bel64] J. S. Bell, *On the Einstein-Podolsky-Rosen paradox*, Physics **1**, 195–200 (1964), Reprinted in J. S. Bell “Speakable and unspeakable in quantum mechanics”, Cambridge University Press, 1987.
- [BN01] F. Benatti and H. Narnhofer, *Additivity of the entanglement of formation*, Phys. Rev. A **63**(4), 042306 (2001), quant-ph/0005126.
- [BNO02] O. Biham, M. A. Nielsen and T. J. Osborne, *Entanglement monotone derived from Grover's algorithm*, Phys. Rev. A **65**, 062312 (2002), quant-ph/0112097.

-
- [Bru02] D. Bruß, *Characterizing entanglement*, J. Math. Phys. **43**(9), 4237–4251 (2002), quant-ph/0110078.
 - [Bur69] D. Bures, *An Extension of Kakutani’s Theorem on Infinite Product Measures to the Tensor Product of Semifinite w^* -Algebras*, Trans. Am. Math. Soc. **135**, 199–212 (1969).
 - [BV04] F. G. S. L. Brandão and R. O. Vianna, *Witnessed Entropy of Entanglement*, (May 2004), quant-ph/0405096.
 - [BW92] C. H. Bennett and S. J. Wiesner, *Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states*, Phys. Rev. Lett. **69**, 2881–2884 (1992).
 - [Coh98] O. Cohen, *Unlocking Hidden Entanglement with Classical Information*, Phys. Rev. Lett. **80**(11), 2493–2496 (1998).
 - [CW04] M. Christandl and A. Winter, *“Squashed Entanglement” – An Additive Entanglement Measure*, J. Math. Phys. **45**(3), 829–840 (2004), quant-ph/0308088.
 - [DCLB00] W. Dür, J. I. Cirac, M. Lewenstein and D. Bruß, *Distillability and partial transposition in bipartite systems*, Phys. Rev. A **61**(6), 062313 (2000), quant-ph/9910022.
 - [DFM⁺99] D. DiVincenzo, C. Fuchs, H. Mabuchi, J. Smolin, A. Thapliyal and A. Uhlmann, *Entanglement of Assistance*, in *Quantum Computing and Quantum Communications: First NASA International Conference, QCQC’98*, edited by C. P. Williams, volume 1509 of *Lecture Notes in Computer Science*, pages 247–257, Springer-Verlag, 1999, quant-ph/980333.
 - [DH99] M. J. Donald and M. Horodecki, *Continuity of relative entropy of entanglement*, Phys. Lett. A **264**, 257–260 (1999), quant-ph/9910002.
 - [DHR02] M. J. Donald, M. Horodecki and O. Rudolph, *The uniqueness theorem for entanglement measures*, J. Math. Phys. **43**(9), 4252–4272 (September 2002), quant-ph/0105017.
 - [DSS⁺00] D. P. DiVincenzo, P. W. Shor, J. A. Smolin, B. M. Terhal and A. V. Thapliyal, *Evidence for bound entangled states with negative partial transpose*, Phys. Rev. A **61**(6), 062312 (2000), quant-ph/9910026.
 - [EAP03] J. Eisert, K. Audenaert and M. B. Plenio, *Remarks on entanglement measures and non-local state distinguishability*, J. Phys. A: Math. Gen. **36**, 5605–5615 (2003), quant-ph/0212007.
 - [Eis03] J. Eisert, *Script of lecture in course: Quantum Information II*, 2003, <http://jense.qip.org/QIT.ps>.

- [Eke91] A. K. Ekert, *Quantum Cryptography Based on Bell's Theorem*, Phys. Rev. Lett. **67**(6), 661–663 (1991).
- [EP03] J. Eisert and M. Plenio, *Introduction to the basics of entanglement theory in continuous-variable systems*, Int. J. Quant. Inf. **1**(4), 479–506 (2003), quant-ph/0312071.
- [EPR35] A. Einstein, B. Podolsky and N. Rosen, *Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?*, Phys. Rev. **47**, 777–780 (1935).
- [ESP02] J. Eisert, C. Simon and M. B. Plenio, *On the quantification of entanglement in infinite-dimensional quantum systems*, J. Phys. A: Math. Gen. **35**, 3911–3923 (2002), quant-ph/0112064.
- [FC95] C. A. Fuchs and C. M. Caves, *Mathematical Techniques for Quantum Communication*, Open Systems & Information Dynamics **3**, 1 (1995), quant-ph/9604001.
- [Gro96] L. Grover, *A Fast Quantum Mechanical Algorithm for Database Search*, in *Proc. 28th Annual ACM Symposium on the Theory of Computation*, pages 212–219, New York, NY, 1996, ACM Press, New York.
- [Gro97] L. K. Grover, *Quantum Mechanics Helps in Searching for a Needle in a Haystack*, Phys. Rev. Lett. **79**, 325–328 (1997), quant-ph/9706033.
- [GWK⁺03] G. Giedke, M. M. Wolf, O. Krueger, R. Werner and J. I. Cirac, *Entanglement of Formation for Symmetric Gaussian States*, Phys. Rev. Lett. **91**, 107901 (2003), quant-ph/0304042.
- [HHH96] M. Horodecki, P. Horodecki and R. Horodecki, *Separability of mixed states: Necessary and sufficient conditions*, Phys. Lett. A **223**(1-2), 1–8 (November 1996), quant-ph/9605038.
- [HHH97] M. Horodecki, P. Horodecki and R. Horodecki, *Inseparable Two Spin- $\frac{1}{2}$ Density Matrices Can Be Distilled to a Singlet Form*, Phys. Rev. Lett. **78**(4), 574–577 (January 1997).
- [HHH98] M. Horodecki, P. Horodecki and R. Horodecki, *Mixed-State Entanglement and Distillation: Is there a “Bound” Entanglement in Nature?*, Phys. Rev. Lett. **80**(24), 5239–5242 (June 1998), quant-ph/9801069.
- [HHH99a] M. Horodecki, P. Horodecki and R. Horodecki, *General teleportation channel, singlet fraction, and quasidistillation*, Phys. Rev. A **60**(3), 1888–1898 (September 1999), quant-ph/9807091.
- [HHH99b] P. Horodecki, M. Horodecki and R. Horodecki, *Bound Entanglement Can Be Activated*, Phys. Rev. Lett. **82**(5), 1056–1059 (1999), quant-ph/9806058.

- [HHH00] M. Horodecki, P. Horodecki and R. Horodecki, *Limits for Entanglement Measures*, Phys. Rev. Lett. **84**(9), 2014–2017 (2000), quant-ph/9908065.
- [HHT01] P. M. Hayden, M. Horodecki and B. Terhal, *The asymptotic entanglement cost of preparing a quantum state*, J. Phys. A: Math. Gen. **34**, 6891–6898 (2001), quant-ph/0008134.
- [Hir02] T. Hiroshima, *An entanglement measure based on the capacity of dense coding*, Phys. Lett. A **301**, 263–268 (2002), quant-ph/0204086.
- [Hor97] P. Horodecki, *Separability criterion and inseparable mixed states with positive partial transposition*, Phys. Lett. A **232**, 333–339 (August 1997), quant-ph/9703004.
- [Hor01] M. Horodecki, *Entanglement Measures*, Quantum Inf. Comput. **1**(1), 3 (2001), <http://www.rintonpress.com/journals/qic-1-1/miarprz3.pdf>.
- [Hüb92] M. Hübner, *Explicit computation of the Bures distance for density matrices*, Phys. Lett. A **163**, 239–242 (1992).
- [HW97] S. Hill and W. Wootters, *Entanglement of a Pair of Quantum Bits*, Phys. Rev. Lett. **78**(26), 5022–5025 (1997), quant-ph/9703041.
- [Key02] M. Keyl, *Fundamentals of quantum information theory*, Physics Reports **369**(5), 431–548 (2002), quant-ph/0202122.
- [KGLC03] B. Kraus, G. Giedke, M. Lewenstein and J. Cirac, *Entanglement properties of Gaussian States*, Fortschr. Phys. **51**(4-5), 305–312 (2003).
- [LCOK03] S. Lee, D. P. Chi, S. D. Oh and J. Kim, *Convex-roof extended negativity as an entanglement measure for bipartite quantum systems*, Phys. Rev. A **68**(6), 062304 (2003), quant-ph/0310027.
- [LP99] H.-K. Lo and S. Popescu, *Classical Communication Cost of Entanglement Manipulation: Is Entanglement an Inverconvertible Resource?*, Phys. Rev. Lett. **83**(7), 1459–1462 (1999), quant-ph/9902045.
- [MSV04] F. Morikoshi, M. F. Santos and V. Vedral, *Accessability of physical states and non-uniqueness of entanglement measure*, J. Phys. A: Math. Gen. **37**(22), 5887–5893 (2004), quant-ph/0306032.
- [Myr04] J. Myrheim, *Separable density matrices*, (unpublished), April 2004.
- [NC00] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, UK, 2000.
- [Nie00] M. A. Nielsen, *Continuity bounds for entanglement*, Phys. Rev. A **61**(6), 064301 (2000), quant-ph/9908086.

- [Nie01] M. A. Nielsen, *On the units of bipartite entanglement: is sixteen ounces of entanglement always equal to one pound?*, J. Phys. A: Math. Gen. **34**, 6987–6995 (2001), quant-ph/0011063.
- [Oza00] M. Ozawa, *Entanglement measures and the Hilbert-Schmidt distance*, Phys. Lett. A **268**, 158–160 (2000), quant-ph/0002036.
- [Per95] A. Peres, *Higher order Schmidt decompositions*, Phys. Lett. A **202**, 16–17 (1995), quant-ph/9504006.
- [Per96] A. Peres, *Separability criterion for density matrices*, Phys. Rev. Lett. **77**(8), 1413–1415 (August 1996), quant-ph/9604005.
- [PR97] S. Popescu and D. Rohrlich, *Thermodynamics and the measure of entanglement*, Phys. Rev. A **56**(5), R3319–R3321 (1997), quant-ph/9610044.
- [PR00] A. O. Pittenger and M. H. Rubin, *Separability and Fourier representations of density matrices*, Phys. Rev. A **62**, 032313 (2000), quant-ph/0001014.
- [Pre98] J. Preskill, Lecture Notes for Physics 229: Quantum Information and Computation, September 1998, <http://www.theory.caltech.edu/people/preskill/ph229/>.
- [RE04] G. Rigolin and C. O. Escobar, *Lower bounds on the entanglement of formation for general Gaussian states*, Phys. Rev. A **69**, 012307 (2004), quant-ph/0307023.
- [Rud01] O. Rudolph, *A new class of entanglement measures*, J. Math. Phys. **42**(11), 5306–5314 (November 2001), math-ph/0005011.
- [San04] E. Santos, *Optical tests of Bell’s inequalities not resting upon the absurd fair sampling assumption*, (2004), quant-ph/0401003.
- [Sha48] C. E. Shannon, *A Mathematical Theory of Communication*, Bell System Technical Journal **27**, 379–423, 623–656 (July, October 1948), <http://cm.bell-labs.com/cm/ms/what/shannonday/paper.html>.
- [Shi02] T. Shimono, *Towards Additivity of Entanglement of Formation*, in *Unconventional Models of Computation: Third International Conference, UMC 2002, Kobe, Japan*, edited by F. P. C. S. Calude, M. J. Dinneen, volume 2509 of *Lecture Notes in Computer Science*, pages 252–263, Springer-Verlag, 2002.
- [Sho94] P. W. Shor, *Algorithms for quantum computation: Discrete logarithms and factoring*, in *Proceedings of the 35th Annual Symposium on the Foundations of Computer Science*, edited by S. Goldwasser, pages 124–134, Los Alamitos, CA, 1994, IEEE Computer Society.
- [Sho97] P. W. Shor, *Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer*, SIAM J. Comput. **26**(5), 1484–1509 (October 1997), quant-ph/9508027.

- [Sim00] R. Simon, *Peres-Horodecki Separability Criterion for Continuous Variable Systems*, Phys. Rev. Lett. **84**(12), 2726–2729 (2000), quant-ph/9909044.
- [SM95] J. Schlienz and G. Mahler, *Description of entanglement*, Phys. Rev. A **52**(6), 4396–4404 (December 1995).
- [SST01] P. W. Shor, J. A. Smolin and B. M. Terhal, *Nonadditivity of Bipartite Distillable Entanglement Follows from a Conjecture on Bound Entangled Werner States*, Phys. Rev. Lett. **86**(12), 2681–2684 (2001), quant-ph/0010054.
- [SW02] B. Schumacher and M. D. Westmoreland, *Relative entropy in quantum information theory*, in *Quantum Computation and Information*, edited by S. J. Lomonaco Jr. and H. E. Brant, volume 305 of *Contemporary Mathematics*, American Mathematical Society, 2002, quant-ph/0004045.
- [TH02] C. H. Thompson and H. Holstein, *The “Chaotic Ball” model: local realism and the Bell test “detection loophole”*, (2002), quant-ph/0210150.
- [Tha99] A. V. Thapliyal, *Multipartite pure-state entanglement*, Phys. Rev. A **59**(5), 3336 (May 1999), quant-ph/9811091.
- [TM71] M. Tribus and E. C. McIrvine, *Energy and information*, Scientific American **225**(3), 179–188 (September 1971).
- [TV00] B. M. Terhal and K. G. H. Vollbrecht, *Entanglement of Formation for Isotropic States*, Phys. Rev. Lett. **85**(12), 2625–2628 (2000), quant-ph/0005062.
- [Uhl76] A. Uhlmann, *The “transition probability” in the state space of a *-algebra*, Rep. Math. Phys. **9**(2), 273–279 (1976).
- [Uhl98] A. Uhlmann, *Entropy and Optimal Decompositions of States Relative to a Maximal Commutative Subalgebra*, Open Systems & Information Dynamics **5**, 209–227 (1998), quant-ph/9704017.
- [Uhl00] A. Uhlmann, *Fidelity and concurrence of conjugated state*, Phys. Rev. A **62**(3), 032307 (2000), quant-ph/9909060.
- [VDC02] G. Vidal, W. Dür and J. I. Cirac, *Entanglement Cost of Bipartite Mixed States*, Phys. Rev. Lett. **89**(2), 027901 (2002), quant-ph/0112131.
- [Ved02] V. Vedral, *The role of relative entropy in quantum information theory*, Rev. Mod. Phys. **74**(1), 197–234 (January 2002), quant-ph/0102094.
- [Vid00] G. Vidal, *Entanglement monotones*, J. Mod. Opt. **47**, 355–376 (2000), quant-ph/9807077.
- [vN32] J. von Neumann, *Mathematische Grundlagen der Quantenmechanik*, Number XXXVIII in Die Grundlehren der Mathematischen Wissenschaften in Einzeldarstellungen, Springer, 1932.

- [vN55] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton University Press, 1955.
- [VP98] V. Vedral and M. Plenio, *Entanglement measures and purification procedures*, Phys. Rev. A **57**(3), 1619–1633 (1998), quant-ph/9707035.
- [VP00] S. Virmani and M. B. Plenio, *Ordering states with entanglement measures*, Phys. Lett. A **268**, 31–34 (2000), quant-ph/001119.
- [VPC04] F. Verstraete, M. Popp and J. Cirac, *Entanglement versus Correlations in Spin Systems*, Phys. Rev. Lett. **92**(2), 027901 (2004), quant-ph/0307009.
- [VPJK97] V. Vedral, M. B. Plenio, K. Jacobs and P. L. Knight, *Statistical inference, distinguishability of quantum states, and quantum entanglement*, Phys. Rev. A **56**(6), 4452–4455 (December 1997), quant-ph/9703025.
- [VPRK97] V. Vedral, M. Plenio, M. Rippin and P. Knight, *Quantifying Entanglement*, Phys. Rev. Lett. **78**(12), 2275–2279 (1997), quant-ph/9702027.
- [VT99] G. Vidal and R. Tarrach, *Robustness of entanglement*, Phys. Rev. A **59**(1), 141–155 (January 1999), quant-ph/9806094.
- [VW01] K. G. H. Vollbrecht and R. F. Werner, *Entanglement measures under symmetry*, Phys. Rev. A **64**(6), 062307 (December 2001), quant-ph/0010095.
- [VW02a] G. Vidal and R. F. Werner, *Computable measure of entanglement*, Phys. Rev. A **65**, 032314 (2002), quant-ph/0102117.
- [VW02b] K. G. H. Vollbrecht and M. M. Wolf, *Activating Distillation with an Infinitesimal Amount of Bound Entanglement*, Phys. Rev. Lett. **88**(24), 247901 (2002), quant-ph/0201103.
- [Wan01] A. M. Wang, *Generalization of the entanglement of formation for multi-party systems*, (2001), quant-ph/0011040 v4.
- [Wer89] R. F. Werner, *Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model*, Phys. Rev. A **40**(8), 4277–4281 (October 1989).
- [WGK⁺04] M. M. Wolf, G. Giedke, O. Krueger, R. Werner and J. Cirac, *Gaussian Entanglement of Formation*, Phys. Rev. A **69**, 052320 (2004), quant-ph/0306177.
- [Whi98] A. Whitaker, *John Bell and the most profound discovery of science*, Physics World **11**(12), 29–34 (December 1998), <http://physicsweb.org/article/world/11/12/8>.
- [Woo98] W. Wootters, *Entanglement of Formation of an Arbitrary State of Two Qubits*, Phys. Rev. Lett. **80**(10), 2245–2248 (1998), quant-ph/9709029.

- [Woo01] W. K. Wootters, *Entanglement of formation and concurrence*, Quantum Inf. Comput. **1**(1), 27–44 (2001), <http://www.rintonpress.com/journals/qic-1-1/eof2.pdf>.
- [WT99] C. Witte and M. Trucks, *A new entanglement measure induced by the Hilbert-Schmidt norm*, Phys. Lett. A **257**, 14–20 (1999), [quant-ph/9811027](http://arxiv.org/abs/quant-ph/9811027).
- [ŻHSL98] K. Życzkowski, P. Horodecki, A. Sanpera and M. Lewenstein, *Volume of the set of separable states*, Phys. Rev. A **58**, 883–892 (August 1998), [quant-ph/9804024](http://arxiv.org/abs/quant-ph/9804024).

Index

- abstract measure, 25
- Alice and Bob, 6
- asymptotic regime, 25
- base norm, 39
- Bell basis, 6
- Bell states, 6
- biorthogonal, 11
- bipartite state, 6
- bit flip operator, 5, 14
- Bloch sphere, 4
- bound entanglement, 23
- Bures metric, 34
- composite system, 5
- concurrence, 36
- convex roof measure, 36
- creation, entanglement of, *see* entanglement, of formation
- density matrix, *see* density operator, 7
- density operator, 7
 - realisation of, 7
 - reduced, 9
- distillable entanglement, *see* entanglement, distillable
- ensemble, 7
- entangled state, 6
- entanglement
 - cost
 - mixed states, 29
 - pure states, 27
 - distillable
 - mixed states, 29
 - pure states, 27
 - distillation, 16–19
 - localizable, 41
 - measure, 25
 - uniqueness theorem, 28
 - of assistance, 40
 - of creation, *see* entanglement, of formation
 - of distillation, *see* entanglement, distillable
 - of formation, 35
 - squashed, 39
- entropy of entanglement, 26
 - relative, 34
 - witnessed, 41
- EPR pair, *see* Bell state
- fidelity, 18
- finite regime, 25
- formation, entanglement of, *see* entanglement, of formation
- free entanglement, 23
- GHZ state, 43
- Local Operations and Classical Communication, *see* LOCC
- LOCC, 14
- LQCC, *see* LOCC
- maximal knowledge, 7
- measurement
 - generalised, 11–13
 - projective, 11
- measurement operators, 11
- mixed state, 7
- multipartite state, 6

- multipartite system, 43
- multiseparable, 44
- negativity, 37
 - logarithmic, 38
- NPPT, 24
- operational measure, 25
- partial trace, 9
- Pauli operators, 5
- Peres-Horodecki criterion, 23
- phase flip operator, 5
- positive operator, 8
- PPT-criterion, 23
- product state, 6, 21
- pure state, 7
- QLCC, *see* LOCC
- qubit, 3
- realisation, 7
- relative entropy, 33
 - of entanglement, 34
- robustness of entanglement, 39
- Schmidt coefficients, 11
- Schmidt decomposition, 10
- Schmidt number, 21
- Schmidt rank, 21
- separability, 21
- separable state, 10, 21
- Shannon entropy, 26
- superadditivity, 41
 - strong, 40
- trace norm, 28, 31, 38
- trace out, 8
- Uhlmann's transition probability, 35
- von Neumann entropy, 26
 - reduced, 26
- X operator, 4, 5
- Y operator, 4, 5
- Z operator, 4, 5